



Rapid evolution of antimicrobial peptide genes in an insect host–social parasite system



Silvio Erler^{a,*}, Patrick Lhomme^b, Pierre Rasmont^b, H. Michael G. Lattorff^{a,c,1}

^a Institut für Biologie, Molekulare Ökologie, Martin-Luther-Universität Halle-Wittenberg, Hoher Weg 4, 06099 Halle (Saale), Germany

^b Laboratoire de Zoologie, Université de Mons, Place du Parc 20, 7000 Mons, Belgium

^c German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

ARTICLE INFO

Article history:

Received 21 October 2013

Received in revised form 8 January 2014

Accepted 7 February 2014

Available online 14 February 2014

Keywords:

Social insect

Co-evolution

Innate immunity

Bumblebee

Bombus

Host–parasite

ABSTRACT

Selection, as a major driver for evolution in host–parasite interactions, may act on two levels; the virulence of the pathogen, and the hosts' defence system. Effectors of the host defence system might evolve faster than other genes e.g. those involved in adaptation to changes in life history or environmental fluctuations. Host–parasite interactions at the level of hosts and their specific social parasites, present a special setting for evolutionarily driven selection, as both share the same environmental conditions and pathogen pressures.

Here, we study the evolution of antimicrobial peptide (AMP) genes, in six host bumblebee and their socially parasitic cuckoo bumblebee species. The selected AMP genes evolved much faster than non-immune genes, but only *defensin-1* showed significant differences between host and social parasite. Nucleotide diversity and codon-by-codon analyses confirmed that purifying selection is the main selective force acting on bumblebee defence genes.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Host–parasite interactions are shaped by a wide range of biotic and abiotic factors. Under natural conditions, variations in parasite (pathogen) load and virulence, as well as variations in host susceptibility and immune responses, control host–parasite dynamics (Anderson and May, 1982; Gandon et al., 2001; Schmid-Hempel and Ebert, 2003). Co-evolutionary arms races are mostly characterized as reciprocal processes of adaptation and counter-adaptation between parasites and hosts (Dawkins and Krebs, 1979). Positive selection of the defence mechanisms clearly reflects the importance of pathogenic organisms in host evolution, because immunity related proteins are functionally important in the host immune system and play an important role in adapting to novel pathogens or pathogen genotypes.

Parallel evolution at the level of amino acid changes are characterized by parallel replacements between at least two related but distinct species possessing a common ancestor. Convergent evolu-

tion describes the same amino acid replacement with the same outcome, in two unrelated species without any common ancestor (Nei and Kumar, 2000). Both, parallel and convergent changes in amino acids are a sign for strong positive selection. However, under natural conditions parallel and convergent evolution have been very rarely observed (Doolittle, 1994).

Evolutionary forces acting on DNA can be characterized by measurable changes of synonymous (d_s) and non-synonymous (d_n) substitution rates in coding regions (Nielsen, 2005; Yang and Bielawski, 2000). The d_n/d_s ratio (ω) classifies possible occurring selection events in three different groups: $\omega > 1$, diversifying selection (positive selection); $\omega = 1$, no selection (neutral evolution) and $\omega < 1$, purifying selection (negative selection) (reviewed in Wagner, 2002).

Innate insect immune systems categorise the majority of parasites into four groups: viruses, gram-positive and gram-negative bacteria, and fungi or yeasts (Hultmark, 1993; Lemaitre and Hoffmann, 2007). Selection may take place separately at genes specific for each group, or globally against one of the groups. Social insects are model organisms for investigating adaptive evolution in the innate immune system, as group living increases their vulnerability to diseases, especially since the group is composed of closely related individuals (Schmid-Hempel, 1998).

Less attention has been paid to host–parasite systems where both host and parasite are closely related, sharing similar life his-

* Corresponding author. Tel.: +49 (0)345 55 26 305; fax: +49 (0)345 55 27 264.

E-mail addresses: erler.silvio@gmail.com (S. Erler), patrick_lhomme@hotmail.fr (P. Lhomme), pierre.rasmont@umons.ac.be (P. Rasmont), michael.lattorff@zoologie.uni-halle.de (H. Michael G. Lattorff).

¹ Present address: Institut für Biologie, Tierphysiologie, Martin-Luther-Universität Halle-Wittenberg, Domplatz 4, 06099 Halle (Saale), Germany.

tory traits. This is known as social parasitism or brood parasitism and is found in both birds and social insects (e.g. ants and bumblebees, reviewed in [Kilner and Langmore, 2011](#)). Host bumblebees and their social parasites, cuckoo bumblebees ([Fisher, 1988](#); [van Honk et al., 1981](#)) share the same, but timeshifted annual life cycle, and corresponding environmental conditions ([Alford, 1975](#); [Sladen, 1912](#)). Host bumblebee colonies are comprised of drones (males), workers and a single bumblebee queen who initiates nest foundation in early spring. During colony development, workers are produced exclusively and only at the end of the season are new sexuals (drones and queens) produced for the forthcoming season ([Alford, 1975](#); [Sladen, 1912](#)). Cuckoo bumblebee females invade host nests in spring, killing the host queen and leaving host workers to take care of the cuckoo female's brood. In contrast to the host, cuckoo females produce only male and queen offspring, lacking a worker caste ([Alford, 1975](#); [Sladen, 1912](#)). This kind of parasitism is assigned to queen-intolerantinquilines (reviewed in [Brandt et al., 2005](#)). Within this asymmetric, inter-specific arms race ([Dawkins and Krebs, 1979](#)), cuckoo bumblebees may be specialist or generalist, being a mono- or multiple-host social parasite, depending on the host species range ([Loken, 1984](#)).

Multiple reports of plant and animal evolutionary adaptations suggest that the environment plays an important role in gene evolution and associated phenotypic shifts (reviewed in [Levasseur et al., 2007](#); [Salamini et al., 2010](#)). Sharing the same environmental conditions (i.e. food source, homeostatic nest condition and symbionts, non-coevolving saprophytes, omnipresent microorganisms at nest and hibernation sites), including co-evolving parasites and pathogens, might force parallel evolution of parasite/pathogen defence mechanisms against common microbes and viruses, in bumblebee hosts and their cuckoo bee parasites. Environmental conditions (e.g., temperature, humidity and light) and shifts in the microhabitat or diets show a strong impact (37.5%; [Fuller et al., 2011](#)) on immunocompetence and pathogen susceptibility in social insects ([Bulmer and Crozier, 2006](#); [Fuller et al., 2011](#)). Transitions to new habitats represent the exposure of the host to novel pathogens, which could direct rapid, adaptive changes in immune proteins ([Bulmer and Crozier, 2006](#)). As a sign of adaptive evolution, genes involved in the immune defences of various plants and animals, typically show a faster rate of nucleotide and amino acid substitutions (non-synonymous), than non-immunity related genes ([McTaggart et al., 2012](#); [Obbard et al., 2009](#); [Sackton et al., 2007](#); [Tiffin and Moeller, 2006](#); [Trowsdale and Parham, 2004](#); reviewed in [Bulmer, 2010](#)).

Positive selection and rapid gene duplication as factors influencing evolution have been demonstrated in social insects for antimicrobial peptides (AMPs) (termicin – [Bulmer and Crozier, 2004](#)), gram-negative bacteria-binding proteins and relish in termites ([Bulmer and Crozier, 2006](#)) and several immune genes in ants ([Viljakainen and Pamilo, 2008](#); [Viljakainen et al., 2009](#)). Positive selection was detected mostly in the mature region of the AMPs, whereas the signal and pro-regions seem to evolve neutrally ([Lazzaro and Clark, 2003](#); [Viljakainen and Pamilo, 2008](#)). For termicin especially, the expressed mature peptide appears to have diverged more rapidly than the 3'UTR ([Bulmer and Crozier, 2004](#)). In addition, a population genetic analysis of nucleotide intra-specific polymorphism and inter-specific divergence indicated that a positive selection driven selective sweep reduced polymorphisms in the AMP termicin ([Bulmer et al., 2010](#)). Hence, if the immune system adapts to parasites/pathogens in similar ways in related species (i.e. host and cuckoo bumblebee species), we would expect to observe congeneric genes experiencing positive selection in different lineages of the same affiliation.

Social insects show a reduced number of immune genes relative to solitary species ([Evans et al., 2006](#)), and so may compensate for the reduction in immunity gene variance through group level 'so-

cial immunity' ([Cremer et al., 2007](#); [Richter et al., 2012](#); [Traniello et al., 2002](#)).

Social parasites and their hosts are frequently very close phylogenetic relatives that might influence the ease of evolutionary adaptations between host and social parasite on both sides ([Davies et al., 1989](#)). Here we tested whether parasite or pathogen driven evolutionary adaptations (parallel evolution of AMP genes) can be observed in closely related host-social parasite couples sharing the same environmental conditions i.e. parasite pressure ([Erler et al., 2012](#)). Six specialised host/social parasite bumblebee couples were used to determine the type and strength of selection on AMP genes, both within and between host and social parasite species.

2. Material and methods

2.1. Bumblebee samples

Bumblebee drones of six bumblebee hosts and their respective cuckoo bumblebee species were sampled in three locations across Europe ([Table 1](#)). Bumblebees have a haplo-diploid sex determination system; therefore the haploid drones provide a highly efficient model system for genetic studies as they present a single allele per locus. At each location, host and social parasite couples were caught during foraging flights and immediately stored in ethanol or at -80°C until further processing. Bumblebee species were identified using the taxonomic key of [Maus \(1994\)](#).

2.2. DNA isolation and target gene amplification

The thorax muscles of three individuals per species were used to isolate genomic DNA using the DNeasy Blood & Tissue Kit tissue protocol (Qiagen, Hilden, Germany). Tissue samples were homogenized, followed by proteinase K (600 mAU/mL) treatment for at least 2 h and final DNA elution was conducted twice in 50 μL AE Buffer. Quality and quantity of DNA was determined via NanoDrop ND-1000 (PepLab, Erlangen, Germany).

AMP (*abaecin*, *defensin-1* and *hymenoptaecin*; all bumblebees) and non-immune reference gene – (*EF-1 alpha*, *arginine kinase*, *rhodopsin*, *PEPCK*; only for *B. perezii*) amplification was performed in a thermocycler, with denaturation at 95°C for 4 min; 35 cycles at 95°C for 40 s; 55°C for 30 s, and 72°C for 2 min 20 s, with final elongation at 72°C for 10 min. Each reaction (10 μL) contained 2.0 mM dNTPs, 0.2 μM of each gene-specific forward and reverse primer ([Table 2](#)), 0.25 U of *peqGOLD* Taq-DNA-polymerase (PepLab, Erlangen, Germany) and 1 μL of extracted genomic DNA.

PCR products were checked for correct amplicon size by automated multicapillary electrophoresis using the QIAxcel System with QIAxcel DNA High Resolution Kit (Qiagen, Hilden, Germany), purified using the QIAquick PCR Purification Kit (Qiagen, Hilden, Germany) or SureClean (Bioline, Luckenwalde, Germany), before 96-well plate MTP sequencing by LGC Genomics (Berlin, Germany), based on traditional Sanger sequencing. All fragments were sequenced overlapping in both directions. *Abaecin*, *defensin-1* and *hymenoptaecin* were successfully amplified in all 12 host and social parasite bumblebee species listed in [Table 1](#). The sequenced regions did not cover the entire coding region of the genes but lacked a few nucleotides of the coding regions in either the 3' or 5' end, or both.

A sequenced region of the 16S rRNA was used to confirm species identification of the cuckoo bumblebee species by comparison with reference sequences from GenBank ([Cameron et al., 2007](#)).

AMP gene sequences for all bumblebee species, non-immune and 16S rRNA gene sequences for *B. perezii* are available on GenBank, under the accession numbers: KC662127–38 (*abaecin*),

Table 1

Bumblebee host and social parasite species overview, including sampling location and year.

	Species (subgenera)	Location	WGS coordinates	Country, region	Year
Host	<i>B. (Bombus) lucorum</i>	Nyer	42°30'N 2°19'E	France, Pyrenees	2007
Parasite	<i>B. (Psithyrus) bohemicus</i>	Nyer	42°31'N 2°17'E	France, Pyrenees	2007
Host	<i>B. (Bombus) terrestris xanthopus</i> ^a	Morosaglia	42°15'N 9°11'E	France, Corsica	2007
Parasite	<i>B. (Psithyrus) perezi</i> ^a	Morosaglia	42°24'N 9°11'E	France, Corsica	2007
Host	<i>B. (Megabombus) ruderatus corsicola</i>	Guaitella	42°25'N 9°25'E	France, Corsica	2008
Parasite	<i>B. (Psithyrus) maxillosus italicus</i> ^b	Haut-Asco	42°14'N 8°33'E	France, Corsica	2008
Host	<i>B. (Bombus) terrestris terrestris</i>	Halle (Saale)	51°29'N 11°58'E	Germany, Saxony-Anhalt	2009
Parasite	<i>B. (Psithyrus) vestalis</i>	Halle (Saale)	51°29'N 11°58'E	Germany, Saxony-Anhalt	2009
Host	<i>B. (Melanobombus) lapidarius</i>	Halle (Saale)	51°29'N 11°58'E	Germany, Saxony-Anhalt	2009
Parasite	<i>B. (Psithyrus) rupestris</i>	Halle (Saale)	51°29'N 11°58'E	Germany, Saxony-Anhalt	2009
Host	<i>B. (Thoracobombus) pascuorum</i>	Halle (Saale)	51°29'N 11°56'E	Germany, Saxony-Anhalt	2009
Parasite	<i>B. (Psithyrus) campestris</i>	Halle (Saale)	51°29'N 11°56'E	Germany, Saxony-Anhalt	2009

^a according to [Lecocq et al. \(2013\)](#).^b recently renamed to *Bombus barbutellus maxillosus* ([Lecocq et al. \(2011\)](#)).**Table 2**

Primer information for the immune and non-immune genes.

Gene	Primer	Sequence (5'–3')	T _m	Reference
<i>Abaecin</i>	A1-F	CCGCCACGACCGGGACAATC	67	This study
	A1-R	GAAACGAAACCGCTGCGAA	60	This study
<i>Defensin-1</i>	def_1529-F	AAACGCAGAAAGACAAAACG	54	This study
	def_2933-R	CGAAACGTTTGTCACAG	57	This study
<i>Hymenoptaecin</i>	H-F	ACTGGCTCTCTTCGATGG	60	This study
	H-R	GAAGCTGGCGAGATTCTG	59	This study
<i>Arginine kinase</i>	ArgK2-F	GACAGCAARTCTCTGCTGAAGAA	62	^a
	ArgK2-R	GGTYTTGGCATCGTTGGTAGATAC	67	^a
<i>EF-1 alpha</i>	F2-ForH	GGRCAYAGAGATTTCATCAAGAAC	62	^b
	F2-RevH2	TTGCAAAGCTTCRKGATGCATT	59	^b
<i>PEPCK</i>	FHv4	TGTATRATAATTCGCAAYTTCAC	56	^c
	RHv	CTGCTGGRGTCTAGATCC	59	^c
<i>Rhodopsin</i>	LWRhF	AATTGCTATTAYGARACNTGGGT	58	^d
	LWRhR	ATATGGAGTCCANGCCATRAACCA	64	^d
16S rRNA	16SWb	CACCTGTTTATCAAAAACAT	50	^e
	874-16SIR	TATAGATAGAAACCAATCTG	50	^f

T_m, melting temperature.^a [Kawakita et al. \(2003\)](#).^b [Hines et al. \(2006\)](#).^c [Cameron et al. \(2007\)](#).^d [Mardulyn and Cameron \(1999\)](#).^e [Downton and Austin \(1994\)](#).^f [Cameron et al. \(1992\)](#).

KC662139–50 (*defensin-1*), KC662151–62 (*hymenoptaecin*), KC662163–67 (*B. perezi* genes).

2.3. Sequence analysis and test for selection in AMP genes

ContigExpress and AlignX, implemented in Vector NTI Advance 10.3.0 (Invitrogen, Carlsbad, CA, USA), were used for sequence alignments, visual inspection and manual editing of the sequences. Exon–intron structure of the AMP genes was verified by sequence comparison with AMP gene sequences of *B. t. terrestris* from GenBank (XM_003394654, XM_003395924, XR_132450) and [Erler](#)

[et al. \(2011\)](#), and of *B. impatiens* (GenBank: XM_003491496, XM_003486302, XM_003494885).

DnaSP v5.10.1 ([Librado and Rozas, 2009](#)) was used to calculate nucleotide diversity (synonymous and non-synonymous sites, including Jukes–Cantor correction ([Jukes and Cantor, 1969](#))) for *defensin-1*, *hymenoptaecin* and non-immune genes (GenBank accession numbers see [Cameron et al., 2007](#)) over all species, for the host species only, social parasite species only and for all pair-wise comparisons within hosts and social parasites. A Mann–Whitney *U* (MWU) test, implemented in STATISTICA 8.0 (StatSoft, Tulsa, OK), was used to test for significant differences in nucleotide diversity

between AMP and non-immune genes and between host and social parasite bumblebee species using all pair-wise comparisons within hosts and within social parasites.

Codon-by-codon analyses of evolutionary changes reveal codon specific adaptations, which indicate either parallel or convergent evolution of AMP genes. Two different methods were applied to test both scenarios (reviewed in [Anisimova and Liberles, 2007](#)).

- (1) Selecton version 2.4 ([Stern et al., 2007](#)) calculates nucleotide changes of each codon to detect positive and purifying selection using a Bayesian inference approach. Default settings for 'Positive selection enabled (M8, $\beta + w > 1$)' were used, including *B. t. terrestris* as the template sequence and a reference phylogenetic tree as the initial tree (for details see 2). Selecton infers site-specific K_a/K_s (also known as d_N/d_S) by computing the expectation of the posterior distribution at each site. After determining the probability for positive selection, the assumption of positive selection will be compared to the null model: no positive selection (M8a, $\beta + w = 1$), using a likelihood ratio test (LRT) ([Yang et al., 2000](#)). If there is no significant difference between M8 and M8a, the hypothesis of positive selection can be rejected and no codon shows any kind of positively selected sites.
- (2) Determination of the ancestral state for each codon position ([Nei and Kumar, 2000](#), Chapter 11.4): If several bumblebee species might share the same codon changes from the same ancestral state, such changes will hint towards parallel evolution. The reconstruction of the ancestral state for codons of AMP genes is done according to a reference phylogenetic tree based on the non-immune gene set. The genes *EF-1 alpha*, *arginine kinase*, *rhodopsin* and *PEPCK* ([Cameron et al., 2007](#)) were used to reconstruct the reference host-social parasite phylogeny. Model selection (tree to use: neighbor-joining tree, statistical method: Maximum Likelihood, substitution type: amino acid), implemented in MEGA v5.1 ([Tamura et al., 2011](#)) was used to determine the best model to reconstruct the bumblebee phylogeny from this data set. For both data sets (AMP and non-immune genes) the Jones–Taylor–Thornton (JTT) model, including bootstrap method with 500 replicates, was selected to be the best substitution model. Finally, the ancestral state was estimated using the non-immune gene tree (user tree) as a template for the AMP gene tree. A phylogenetic tree was established including all possible changes at each ancestral state and a list of amino acid changes along the target sequence. Comparing amino acid changes over host and social parasite couples, or only within the group host and within the group social parasite, shows candidates for selection and parallel evolution. A similar function for the analysis of the ancestral state is implemented in Selecton v2.4 and was used to double-check the results from MEGA analysis.

Classical phylogenetic reconstruction based on nucleotide substitutions was used to check for the overall pattern of changes in the AMP gene coding sequences and clustering of host and social parasite species compared to the reference tree (non-immune genes). Model selection revealed the Kimura 2-parameter model, which considers transitional and transversional substitution rates, as the best model. Bootstrap resampling method (1000 replications) was used to verify the topology of the inferred phylogenetic tree.

In addition, to check if evolutionary changes might also occur in non-coding AMP sequences, we analyzed the phylogenetic relationships of the six host and social parasite bumblebee couples using Maximum Likelihood analysis of the genetic information from all three AMP genes (*abaecin*, *defensin-1*, *hymenoptaecin*). Dif-

ferent models were used for: (1) all coding and non-coding regions (model: Tamura 3-Parameter including gamma distribution), (2) only coding regions (model: Kimura 2-Parameter) and (3) only non-coding regions (model: Tamura 3-Parameter), including bootstrap testing of the robustness of the final tree (1000 replications). Different models were selected on the lowest BIC (Bayesian Information Criterion) score using Model selection implemented in MEGA v5.1 ([Tamura et al., 2011](#)).

3. Results

3.1. Evolutionary speed of AMP genes

The sequenced regions did not cover the entire coding region of the genes, lacking a few coding regions in either the 3' or 5' end, or both. In detail, 33 bp of the *abaecin* coding region and up to 965 bp non-coding sequences were successfully amplified for all bumblebee species. *B. pascuorum* (893 bp), followed by *B. ruderatus* (946 bp), showed the lowest nucleotide number for the non-coding sequence due to many species specific deletions. An intron yet to be described was detected in all bumblebee species, located between amino acid seven and eight, with characteristic acceptor and donor splice sites GT/AG. For *defensin-1*, 60 bp of the pro region and 120 bp of the mature peptide coding region were used for analysis and up to 1031 bp non-coding sequences (*B. campestris* showed the lowest number – 418 bp). Two bumblebee species had large non-coding region sequence gaps for *defensin-1* (*B. m. italicus*: ca. 180 bp, *B. perezii*: ca. 600 bp) due to unsatisfactory sequencing results. Finally, 21 bp of the pro region and 210 bp of the *hymenoptaecin* mature peptide sequence were sequenced, including up to 900 bp non-coding sequence. Once again, *B. campestris* had the shortest non-coding sequence (432 bp) because of many species specific deletions, followed by *B. lapidarius*, *B. r. corsicola* and *B. pascuorum*. Details on the pro and mature peptide region and the non-coding introns of each AMP, over all bumblebee species, are visualized in the [Supplementary Figs. S1–S3](#), using nucleotide sequence alignments (using MAFFT version 6; [Katoh and Toh, 2010](#); [Katoh et al., 2002](#)) and the AMP annotation of *B. ignitus* ([Choi et al., 2008](#)).

The coding regions of all analyzed antimicrobial peptide genes (*abaecin*, *defensin-1*, *hymenoptaecin*) are highly conserved over the 12 bumblebee species. *Defensin-1* homology values (percent identity at nucleotide level) ranged between 94.35% (between *B. campestris* and the group of *B. t. terrestris*, *B. t. xanthopus*, *B. lucorum*) and two groups with 100% (1) *B. t. terrestris*, *B. t. xanthopus*, *B. lucorum* and (2) *B. m. italicus*, *B. perezii*, *B. vestalis*. *Hymenoptaecin* showed nucleotide conservation within a range of 93.45% (between *B. pascuorum* and the group of *B. t. terrestris*, *B. t. xanthopus*, *B. lucorum*) and 100% (between *B. t. terrestris*, *B. t. xanthopus* and *B. lucorum*). *Abaecin* coding region revealed identity ranges between 93% and 100%, but without any distinct clustering due to the small amount of analyzed nucleotides. Only *B. lapidarius* and *B. r. corsicola* showed sequence differences compared with the others in the *abaecin* coding region.

All tested AMP genes evolved much faster than the non-immune genes, irrespective of species affiliation (overall: MWU, $Z = -13.715$, $N = 366$, $P < 0.001$; only host species: MWU, $Z = -7.015$, $N = 78$, $P < 0.001$; only social parasite species: MWU, $Z = -3.160$, $N = 72$, $P = 0.002$). Substitution rates (d_N/d_S) of AMP genes ranged between 0.099–0.672, and 0–0.06 for the non-immune genes ([Table 3](#)). When comparing evolutionary changes between host and social parasite bumblebee species, only *defensin-1* showed significant differences (MWU test, $P = 0.01$, [Table 3](#)) with host species having evolved much faster than the social parasite species.

Table 3

d_N/d_S ratios for host and social parasite immune (*defensin-1*, *hymenoptaecin*) and non-immune genes (*arginine kinase*, *EF-1 alpha*, *PEPCK*, *rhodopsin*). Mann–Whitney *U* (MWU) test shows significant differences between groups ($P < 0.05$).

Gene	d_N/d_S (Overall)	d_N/d_S (Host)	d_N/d_S (Parasite)	MWU-Test (Host vs. Parasite)
<i>Defensin-1</i>	0.243	0.267	0.099	0.001
<i>Hymenoptaecin</i>	0.367	0.225	0.672	0.085
Mean	0.305	0.246	0.386	
<i>Arginine kinase</i>	0.01	0.019	0	0.108
<i>EF-1 alpha</i>	0.005	0.008	0	0.108
<i>PEPCK</i>	0.017	0.043	0	0.002
<i>Rhodopsin</i>	0.035	0.060	0	<0.001
Mean	0.017	0.033	0	

3.2. Non-coding sequences

Comparing nucleotide substitutions of the AMP non-coding sequences (only introns available) revealed no clear signal for either parallel or convergent evolution between hosts and their social parasites for each of the six bumblebee couples. Mostly constant patterns of indels and nucleotide substitutions were observed, demonstrating a similar evolutionary pattern within the group of hosts and social parasites, respectively (see [Supplementary Figs. S1–S3](#)). *Abaecin* showed constant patterns of indels and substitutions between *B. lucorum*, *B. t. terrestris*, and *B. t. xanthopus* (including one obvious insertion – GTAT) and between *B. m. italicus*, *B. perezii*, and *B. vestalis*. The same phenomenon was detectable for *defensin-1* and *hymenoptaecin*, showing constant patterns of indels and substitutions between *B. lucorum*, *B. t. terrestris*, *B. t. xanthopus* (e.g. GCGACTATTCG insertion for *defensin-1*). In addition, reconstructing the phylogenetic relationship of the 12 bumblebee species using non-coding sequence substitution patterns revealed an equal separation in two major groups: the first including the so-

cial parasites and second the host bumblebee group ([Supplementary Fig. S4](#)). Thus, the same pattern was ascertained as in [Fig. 1](#) and also in agreement with the bumblebee phylogeny by [Cameron et al. \(2007\)](#).

3.3. Selection and evolution of AMP genes

In order to investigate the role of positive selection on the evolution of AMP genes, a codon-based Bayesian inference approach was applied. *Hymenoptaecin* showed a significant difference when comparing the two models among the 12 different bumblebee species (log-likelihood M8 model = −361.669, M8a model = −365.744; LRT $P = 0.01$) ([Fig. 2](#)), which may indicate positive selection acting on *hymenoptaecin*. *Defensin-1* did not pass the level of significance (log-likelihood M8 model = −336.601, M8a model = −337.762; LRT $P > 0.05$). Purifying selected sites were detectable for both *Ka/Ks* scores below 1 ([Fig. 2](#)), with a slightly higher amount for *hymenoptaecin* compared to *defensin-1*. Only a minority of all selectable sites (<15%) had an evolutionary speed up to ~5 times than the basic level of the whole gene.

The reconstruction of the ancestral state sequences for *defensin-1* and *hymenoptaecin* did not show any general pattern of parallel evolution between bumblebee host-social parasite couples ([Table 4](#)). Mostly *B. rupestris*, *B. bohemicus* and *B. campestris* – subgenera *Psithyrus* (*Hymenoptaecin* H8Y) – but without a specific common ancestor ([Fig. 1](#)); and *B. t. terrestris*, *B. t. xanthopus*, *B. lucorum* – subgenera *Bombus* – build up a separate group by *Hymenoptaecin* changes Q18D and Q20K, respectively. Synonymous changes along both sequences confirmed this pattern and also include the cluster *B. m. italicus*, *B. perezii* and *B. vestalis* – subgenera *Psithyrus*. Comparing host and social parasite species changes, only codon 10 of *defensin-1* showed an unambiguous separation of the host from the social parasite species by G10A ([Table 4](#)). Additional comparison of nucleotide changes along the AMP and non-immune genes




Fig. 1. Phylogenetic relationships of the six host and social parasite bumblebee couples for the coding sequences of *defensin-1* (A), *hymenoptaecin* (B) and non-immune reference genes (C). Maximum Likelihood analysis was used with Kimura 2-parameter model method (including test of phylogeny: bootstrap with 1000 replications). Note different scaling between A, B and C.




Fig. 2. K_a/K_s scores obtained for each codon position of *hymenoptaecin*. Bayesian models, which assume a statistical distribution to account for heterogeneous K_a/K_s values among sites, are plotted for each codon (M8 model – solid line, allows for positive selection; M8a model – dashed line, only neutral and purifying selection are allowed). (K_a , non-synonymous rate; K_s , synonymous rate at each codon site.)

Table 4

Amino acid substitutions related to the ancestral state for Defensin-1 and Hymenoptaecin.

	Defensin-1						Hymenoptaecin							
Position	5	6	8	10	16	55	2	8	15	18	20	37	46	56
Ancestral	P	L	H	G	D	I	S	H	L	Q	Q	H	V	I
<i>B. lucorum</i>	L	.	Q	.	.	V	.	.	.	D	K	.	.	.
<i>B. bohemicus</i>	.	.	.	A	.	.	.	Y	V	.	.	.	.	.
<i>B. t. xanthopus</i>	L	.	Q	.	.	V	.	.	.	D	K	.	.	.
<i>B. perezi</i>	.	.	.	A	.	.	.	.	V	.	.	.	.	.
<i>B. r. corsicola</i>	.	.	.	D	V	.	.	.	.	.	.	.	.	V
<i>B. m. italicus</i>	.	.	.	A	.	.	.	.	V	.	.	.	.	.
<i>B. t. terrestris</i>	L	.	Q	.	.	V	.	.	.	D	K	.	.	.
<i>B. vestalis</i>	.	.	.	A	.	.	.	.	V	.	.	.	.	.
<i>B. lapidarius</i>	.	.	.	D	.	V	.	.	.	.	.	.	M	.
<i>B. rupestris</i>	.	F	.	A	.	.	.	Y	V	.	.	R	.	.
<i>B. pascuorum</i>	.	.	.	.	.	.	Y	.	V	.	.	.	.	.
<i>B. campestris</i>	.	.	.	A	.	.	.	Y	I	.	.	.	.	.

also confirmed the absence of a general sign for parallel evolution between host and social parasite couples – AMP genes evolve in a similar pattern but more within bumblebee lineage or subspecies specific than adapted to environmental changes between the host and its social parasites. Fig. 1 illustrates the evident clustering of host and social parasite species into two separate groups for *defensin-1* and *hymenoptaecin* in comparison to the four non-immune genes. Grouping in host and social parasite group was almost consistent between the three different trees, except for the *hymenoptaecin* tree where *B. pascuorum* was placed within the social parasite section (Fig. 1B), which is not surprisingly as *B. pascuorum* (*Thoracobombus*) is the closest relative of the subgenus *Psithyrus* in our dataset (according to Cameron et al., 2007). However, comparing the bootstrap consensus tree with the original tree in respect of bootstrap values (39 for bootstrap consensus tree vs. 5 for original tree), *B. pascuorum* definitely belongs to the host bumblebee group, consistent with any other tree analyzed in this study (see Fig. 1 and Supplementary Fig. S4). The topology of the AMP gene trees at the level of the subgenera (Fig. 1) was in agreement with the extensive bumblebee phylogeny constructed by Cameron et al. (2007). Both clusters, separated in host and social parasite bumblebees, confirmed the absence of parallel evolution between hosts and social parasites sharing the same environment and indicated a strongly similar evolutionary pattern between closely related subspecies. Furthermore, they cluster in accordance with their phylogenetical

information for the major subgenera *Bombus*, *Psithyrus* and *Thoracobombus*. Only *B. lapidarius* and *B. r. corsicola* move from one group to another according to the sequence considered.

4. Discussion

4.1. Evolutionary speed of AMP genes

Bumblebee substitution rates for AMP genes do not differ from other social insects, such as ants and honey bees (*defensin-1*: 0.243 vs. 0.5 in ants, 0.272 in honey bees; *hymenoptaecin*: 0.367 vs. 0.299 in honey bees and *abaecin*: 0.3 vs. 0.3 in ants – Erler and Lattorff, unpublished data; Viljakainen and Pamilo, 2008), indicating high evolutionary rates at the protein level. The median substitution ratio (d_N/d_S) was substantially higher in social insects (bumblebees – 0.305, this study; honey bees – 0.269; ants – 0.500) compared to non-social insects (*Drosophila* – 0.08; *Nasonia* (only *defensin-1* available) – 0.08) (Gao and Zhu, 2010; Sackton et al., 2007; Viljakainen and Pamilo, 2008). Though, substitution ratios of immune genes can be influenced by the group of immune gene (recognition, signalling or effector genes) and the type of outgroup used, (genus or order specific, e.g. *D. simulans*: immunity 0.268, non-immunity 0.082; *D. melanogaster*: immunity 0.207, non-immunity 0.172; outgroup: *D. yakuba*; Schlenke and Begun, 2003). When comparing the

d_N/d_S ratio of non-immune genes, the overall ratio was much lower in bumblebees (0.017) than in honey bees (0.036) and flies (0.045) (Viljakainen and Pamilo, 2008), which may be explained by the low number of non-immune genes.

A McDonald–Kreitman test (McDonald and Kreitman, 1991); which compares non-synonymous and synonymous changes and contrasts within-species polymorphism to fixed differences between species; could not be used to test for within population differences in adaptation due to the limited low sample size for the majority of rare parasitic cuckoo bumblebee species. However it should be noted that estimates of the rate of adaptive substitution can be influenced by factors such as population demography (Hughes, 2007).

4.2. Selection and evolution of AMP genes

Immune systems of social insects (e.g. ants, bees and termites) and dipteran insects (e.g. flies and mosquitoes) may respond differently to the selection pressure caused by microbial parasites and pathogens. No evidence for positive selection has been found in antibacterial or antifungal peptide genes of *Drosophila* and *Bombyx mori* (Jiggins and Kim, 2005; Lazzaro and Clark, 2003; Sackton et al., 2007; Yang et al., 2011) and immune genes, including several AMPs of *Anopheles* (Lehmann et al., 2009; Parmakelis et al., 2008; Simard et al., 2007) indicated no sign for co-evolution with parasites and pathogens. Host–parasite arms races may involve strong selection, but only on a relatively small subset of the immune system, such as Imd- and RNAi-pathway genes (Obbard et al., 2009).

Insects might differ in how they respond to parasite pressure, as positive selection acting on AMP genes has so far been found only in social insect species. Further evidence was found in this study, in the form of positive selection acting on *hymenoptaecin*. Yet irrespective of selection pressure, qualities of AMPs influence important host parameters, such as survivorship after bacterial infection (Coggins et al., 2012). As no signal for positive selection has been shown in AMPs for honeybees, except for *defensin-2*, (Harpur and Zayed, 2013) and in only a few ant, bumblebee and termite genes; immune genes may possibly attack a wide spectrum of non-coevolving bacteria (saprophytes), or selection is acting instead for speed and efficiency of AMP production (Sackton et al., 2007; Simard et al., 2007). Positive selection may play a limited role in the evolution of innate immune genes and relaxed purifying selection, including high rates of non-synonymous polymorphisms and divergence act as central mechanism on immune gene evolution (Harpur and Zayed, 2013). Adaptation of AMPs to non-coevolving, universal saprophytes might be the reason for the absence of any parallel substitution pattern related to the couples of bumblebee host and social parasite species. Purifying selection seems to be the common mode of selected changes among AMP genes of insects, as shown for *defensin-1*, regardless of whether they are social or non-social. Certainly, the above mentioned results of ant and termite studies should be treated with caution, as suggested by Hughes (2012), since the statistical methods used are known to have a very high rate of false positives and no experimental evidence has been provided concerning the biological function of selection affecting amino acid replacements.

The major difference between ants, termites and bumblebees is the life cycle itself. Compared with bumblebees, ants and termites have long-lived colonies, with overlapping generations staying in the same nest, which might create a stable and long-lasting association between the host and its specific pathogens. Such an association might be favourable for positive selection on immunity related genes (Viljakainen and Pamilo, 2008). The evidence of positive selection in AMPs might suggest that AMPs in bumblebees are involved in lineage-specific host–parasite arms races; and no evidence was detected that closely related species experiencing com-

parable selection pressures, adapt to such selection pressures in similar ways.

Several theories were put forward to explain positive selection and AMP gene evolution in social insects and three, mutually non-exclusive hypotheses, have been put forward:

- (1) The rate of non-synonymous substitutions might be elevated in social insects not only because of positive selection, but also by low effective population sizes (see Erler and Latortoff, 2010) and nearly neutral mutations might behave as if neutral (Bromham and Leys, 2005). A small effective population size would increase the rate of slightly harmful amino acid substitutions in all genes and raise the d_N/d_S ratios (Bromham and Leys, 2005; Viljakainen et al., 2009).
- (2) Whilst purifying selection on innate immunity might be relaxed, and amino acid changes could be allowed if ‘social immunity’ (Cremer et al., 2007) compensates this effect, positive selection driven repeated amino acid replacements at selected sites need to be detected more often if the high parasite pressure is especially severe in social insects (Viljakainen et al., 2009).
- (3) Finally, some AMPs might attack their targets in such a way that evolving resistance is not possible without coordinated changes at many microbial genes. Selection on AMPs would primarily occur when hosts enter new niches and are forced to adapt to novel pathogen species not previously encountered (Tennesen, 2005).

4.3. Conclusion

Host and social parasite bumblebee immune system genes undergo more purifying selection, but also positive selection, than non-immune system genes. The combination of social defence mechanisms (‘social immunity’) and the evolution of physiological defence mechanisms form the current model of a combined social insect defence system against parasites and pathogens, which might compensate for the low number of immune genes and isoforms of AMPs. Additional (social) insect genomes, including comparative studies of immune system related gene evolution, will be needed to finally understand host–parasite mediated defence adaptations and selection occurring on host immune systems.

Acknowledgements

We would like to thank D. Kleber and S. Helbing for the help with lab work and data analysis, and the Bumblebee Genome Consortium (<http://hymenoptera-genome.org/beebase/>) for providing genomic resources. We are grateful to Beth Nicholls for critical linguistic suggestions and helpful comments. Financial support was granted by the Federal Ministry of Education and Research (BMBF) program FUGATO-Plus (FKZ: 0315126 to H.M.G.L.) and by the Fonds pour la Recherche dans l'Industrie et l'Agriculture (FRIA).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.meegid.2014.02.002>.

References

- Alford, D.V., 1975. Bumblebees. Davis-Poynter, London.
- Anderson, R.M., May, R.M., 1982. Coevolution of hosts and parasites. *Parasitology* 85, 411–426.
- Anisimova, M., Liberles, D.A., 2007. The quest for natural selection in the age of comparative genomics. *Heredity* 99, 567–579.

- Brandt, M., Foitzik, S., Fischer-Blass, B., Heinze, J., 2005. The coevolutionary dynamics of obligate ant social parasite systems – between prudence and antagonism. *Biol. Rev.* 80, 251–267.
- Bulmer, M.S., 2010. Evolution of immune proteins in insects. *Encyclopedia of Life Sciences*. John Wiley & Sons, Ltd., Chichester. <http://dx.doi.org/10.1002/9780470015902.a0022889> (<http://www.els.net/>).
- Bulmer, M.S., Crozier, R.H., 2004. Duplication and diversifying selection among termite antifungal peptides. *Mol. Biol. Evol.* 21, 2256–2264.
- Bulmer, M.S., Crozier, R.H., 2006. Variation in positive selection in termite GNBPs and Relish. *Mol. Biol. Evol.* 23, 317–326.
- Bulmer, M.S., Lay, F., Hamilton, C., 2010. Adaptive evolution in subterranean termite antifungal peptides. *Insect Mol. Biol.* 19, 669–674.
- Bromham, L., Leys, R., 2005. Sociality and the rate of molecular evolution. *Mol. Biol. Evol.* 22, 1393–1402.
- Cameron, S.A., Hines, H.M., Williams, P.H., 2007. A comprehensive phylogeny of the bumble bees (*Bombus*). *Biol. J. Linn. Soc.* 91, 161–188.
- Cameron, S.A., Derr, J.N., Austin, A.D., Wooley, J.B., Wharton, R.A., 1992. The application of nucleotide sequence data to phylogeny of the Hymenoptera: a review. *J. Hym. Res.* 1, 63–79.
- Choi, Y.S., Choo, Y.M., Lee, K.S., Yoon, H.J., Kim, I., Je, Y.H., Sohn, H.D., Jin, B.R., 2008. Cloning and expression profiling of four antibacterial peptide genes from the bumblebee *Bombus ignitus*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 150, 141–146.
- Coggins, S.A., Estévez-Lao, T.Y., Hillyer, J.F., 2012. Increased survivorship following bacterial infection by the mosquito *Aedes aegypti* as compared to *Anopheles gambiae* correlates with increased transcriptional induction of antimicrobial peptides. *Dev. Comp. Immunol.* 37, 390–401.
- Cremer, S., Armitage, S.A., Schmid-Hempel, P., 2007. Social immunity. *Curr. Biol.* 17, R693–R702.
- Davies, N.B., Bourke, A.F., de L Brooke, M., 1989. Cuckoos and parasitic ants: interspecific brood parasitism as an evolutionary arms race. *Trends Ecol. Evol.* 4, 274–278.
- Dawkins, R., Krebs, J.R., 1979. Arms races between and within species. *Proc. Roy. Soc. B Biol. Sci.* 205, 489–511.
- Doolittle, R.F., 1994. Convergent evolution: the need to be explicit. *Trends Biochem. Sci.* 19, 15–18.
- Dowton, M., Austin, A.D., 1994. Molecular phylogeny of the insect order Hymenoptera: apocritan relationships. *Proc. Natl. Acad. Sci. U.S.A.* 91, 9911–9915.
- Erler, S., Lattorff, H.M.G., 2010. The degree of parasitism of the bumblebee (*Bombus terrestris*) by cuckoo bumblebees (*Bombus (Psithyrus) vestalis*). *Insectes Soc.* 57, 371–377.
- Erler, S., Popp, M., Lattorff, H.M.G., 2011. Dynamics of immune system gene expression upon bacterial challenge and wounding in a social insect (*Bombus terrestris*). *PLoS One* 6, e18126.
- Erler, S., Popp, M., Wolf, S., Lattorff, H.M.G., 2012. Sex, horizontal transmission, and multiple hosts prevent local adaptation of *Crithidia bombi*, a parasite of bumblebees (*Bombus* spp.). *Ecol. Evol.* 2, 930–940.
- Evans, J.D., Aronstein, K., Chen, Y.P., Hetru, C., Imler, J.L., Jiang, H., Kanost, M., Thompson, G.J., Zou, Z., Hultmark, D., 2006. Immune pathways and defence mechanisms in honey bees *Apis mellifera*. *Insect Mol. Biol.* 15, 645–656.
- Fisher, R.M., 1988. Observations on the behaviours of three European cuckoo bumble bee species (*Psithyrus*). *Insectes Soc.* 35, 341–354.
- Fuller, C.A., Postava-Davignon, M.A., West, A., Rosengaus, R.B., 2011. Environmental conditions and their impact on immunocompetence and pathogen susceptibility of the Caribbean termite *Nasutitermes acajutlae*. *Ecol. Entomol.* 36, 459–470.
- Gandon, S., Jansen, V.A., van Baalen, M., 2001. Host life history and the evolution of parasite virulence. *Evolution* 55, 1056–1062.
- Gao, B., Zhu, S., 2010. Identification and characterization of the parasitic wasp *Nasonia defensus*: positive selection targeting the functional region? *Dev. Comp. Immunol.* 34, 659–668.
- Harpur, B.A., Zayed, A., 2013. Accelerated evolution of innate immunity proteins in social insects: adaptive evolution or relaxed constraint? *Mol. Biol. Evol.* 30, 1665–1674.
- Hines, H.M., Cameron, S.A., Williams, P.H., 2006. Molecular phylogeny of the bumble bee subgenus *Pyrobombus* (Hymenoptera: Apidae: *Bombus*) with insights into gene utility for lower level analysis. *Invertebr. Syst.* 20, 289–303.
- Hughes, A.L., 2007. Looking for Darwin in all the wrong places: the misguided quest for positive selection at the nucleotide sequence level. *Heredity* 99, 364–373.
- Hughes, A.L., 2012. Evolution of the β GRP/GNBP/ β -1,3-glucanase family of insects. *Immunogenetics* 64, 549–558.
- Hultmark, D., 1993. Immune reactions in *Drosophila* and other insects: a model for innate immunity. *Trends Genet.* 9, 178–183.
- Jiggins, F.M., Kim, K.W., 2005. The evolution of antifungal peptides in *Drosophila*. *Genetics* 171, 1847–1859.
- Jukes, T.H., Cantor, C.R., 1969. Evolution of protein molecules. In: Munro, H.N. (Ed.), *Mammalian Protein Metabolism*. Academic Press, New York, pp. 21–132.
- Katoh, K., Toh, H., 2010. Parallelization of the MAFFT multiple sequence alignment program. *Bioinformatics* 26, 1899–1900.
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30, 3059–3066.
- Kawakita, A., Sota, T., Ascher, J.S., Ito, M., Tanaka, H., Kato, M., 2003. Evolution and phylogenetic utility of alignment gaps within intron sequences of three nuclear genes in bumble bees (*Bombus*). *Mol. Biol. Evol.* 20, 87–92.
- Kilner, R.M., Langmore, N.E., 2011. Cuckoos versus hosts in insects and birds: adaptations, counter-adaptations and outcomes. *Biol. Rev.* 86, 836–852.
- Lazzaro, B.P., Clark, A.G., 2003. Molecular population genetics of inducible antibacterial peptide genes in *Drosophila melanogaster*. *Mol. Biol. Evol.* 20, 914–923.
- Lecocq, T., Lhomme, P., Michez, D., Dellicour, S., Valterová, I., Rasmont, P., 2011. Molecular and chemical characters to evaluate species status of two cuckoo bumblebees: *Bombus barbutellus* and *Bombus maxillosus* (Hymenoptera, Apidae, Bombini). *Sys. Ent.* 36, 453–469.
- Lecocq, T., Vereecken, N.J., Michez, D., Dellicour, S., Lhomme, P., Valterová, I., Rasplus, J.Y., Rasmont, P., 2013. Patterns of genetic and reproductive traits differentiation in Mainland vs. Corsican populations of bumblebees. *PLoS One* 8, e65642.
- Lehmann, T., Hume, J.C., Licht, M., Burns, C.S., Wollenberg, K., Simard, F., Ribeiro, J.M., 2009. Molecular evolution of immune genes in the malaria mosquito *Anopheles gambiae*. *PLoS One* 4, e4549.
- Lemaitre, B., Hoffmann, J., 2007. The host defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* 25, 697–743.
- Levasseur, A., Orlando, L., Bailly, X., Milinkovitch, M.C., Danchin, E.G., Pontarotti, P., 2007. Conceptual bases for quantifying the role of the environment on gene evolution: the participation of positive selection and neutral evolution. *Biol. Rev.* 82, 551–572.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452.
- Loken, A., 1984. Scandinavian species of the genus *Psithyrus* Lepeletier (Hymenoptera: Apidae). *Entomol. Scand. Suppl.* 23, 45.
- Mardulyn, P., Cameron, S.A., 1999. The major opsin in bees (Insecta: Hymenoptera): a promising nuclear gene for higher level phylogenetics. *Mol. Phylogenet. Evol.* 12, 168–176.
- Mauss, V., 1994. Bestimmungsschlüssel für Hummeln. Deutscher Jugendbund für Naturbeobachtungen (DJN), Hamburg.
- McDonald, J.H., Kreitman, M., 1991. Adaptive protein evolution at the Adh locus in *Drosophila*. *Nature* 351, 652–654.
- McTaggart, S.J., Obbard, D.J., Conlon, C., Little, T.J., 2012. Immune genes undergo more adaptive evolution than non-immune system genes in *Daphnia pulex*. *BMC Evol. Biol.* 12, 63.
- Nei, M., Kumar, S., 2000. *Molecular Evolution and Phylogenetics*. Oxford University Press, Oxford.
- Nielsen, R., 2005. Molecular signatures of natural selection. *Annu. Rev. Genet.* 39, 197–218.
- Obbard, D.J., Welch, J.J., Kim, K.-W., Jiggins, F.M., 2009. Quantifying adaptive evolution in the *Drosophila* immune system. *PLoS Genet.* 5, e1000698.
- Parmakelis, A., Slotman, M.A., Marshall, J.C., Awono-Ambene, P.H., Antonio-Nkondjio, C., Simard, F., Caccone, A., Powell, J.R., 2008. The molecular evolution of four anti-malarial immune genes in the *Anopheles gambiae* species complex. *BMC Evol. Biol.* 8, 79.
- Richter, J., Helbing, S., Erler, S., Lattorff, H.G.M., 2012. Social context-dependent immune gene expression in bumblebees (*Bombus terrestris*). *Behav. Ecol. Sociobiol.* 66, 791–796.
- Sackton, T.B., Lazzaro, B.P., Schlenke, T.A., Evans, J.D., Hultmark, D., Clark, A.G., 2007. Dynamic evolution of the innate immune system in *Drosophila*. *Nat. Genet.* 39, 1461–1468.
- Salamin, N., Wüest, R.O., Laverne, S., Thuiller, W., Pearman, P.B., 2010. Assessing rapid evolution in a changing environment. *Trends Ecol. Evol.* 25, 692–698.
- Schlenke, T.A., Begun, D.J., 2003. Natural selection drives *Drosophila* immune system evolution. *Genetics* 164, 1471–1480.
- Schmid-Hempel, P., 1998. *Parasites in Social Insects*. Princeton University Press, Princeton.
- Schmid-Hempel, P., Ebert, D., 2003. On the evolutionary ecology of specific immune defence. *Trends Ecol. Evol.* 18, 27–32.
- Simard, F., Licht, M., Besansky, N.J., Lehmann, T., 2007. Polymorphism at the defensin gene in the *Anopheles gambiae* complex: testing different selection hypotheses. *Infect. Genet. Evol.* 7, 285–292.
- Sladen, F.W.L., 1912. *The humble-bee – Its life-history and how to domesticate it*. Macmillan Co., Limited, London, London.
- Stern, A., Doron-Faigenboim, A., Erez, E., Martz, E., Bacharach, E., Pupko, T., 2007. Selecton 2007: advanced models for detecting positive and purifying selection using a Bayesian inference approach. *Nucleic Acids Res.* 35, W506–W511.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Tennessen, J.A., 2005. Molecular evolution of animal antimicrobial peptides: widespread moderate positive selection. *J. Evol. Biol.* 18, 1387–1394.
- Tiffin, P., Moeller, D.A., 2006. Molecular evolution of plant immune system genes. *Trends Genet.* 22, 662–670.
- Traniello, J.F.A., Rosengaus, R.B., Savoie, K., 2002. The development of immunity in a social insect: evidence for the group facilitation of disease resistance. *Proc. Natl. Acad. Sci. U.S.A.* 99, 6838–6842.
- Trowsdale, J., Parham, P., 2004. Mini-review: defense strategies and immunity-related genes. *Eur. J. Immunol.* 34, 7–17.
- van Honk, C., Röseler, P.F., Velthuis, H.H.W., Malotiaux, M., 1981. The conquest of a *Bombus terrestris* colony by a *Psithyrus vestalis* female. *Apidologie* 12, 57–67.
- Viljakainen, L., Pamilo, P., 2008. Selection on an antimicrobial peptide defensin in ants. *J. Mol. Evol.* 67, 643–652.

- Viljakainen, L., Evans, J.D., Hasselmann, M., Rueppell, O., Tingek, S., Pamilo, P., 2009. Rapid evolution of immune proteins in social insects. *Mol. Biol. Evol.* 26, 1791–1801.
- Wagner, A., 2002. Selection and gene duplication: a view from the genome. *Genome Biol.* 3, reviews1012.1–reviews1012.3.
- Yang, Z., Bielawski, J.P., 2000. Statistical methods for detecting molecular adaptation. *Trends Ecol. Evol.* 15, 496–503.
- Yang, Z., Nielsen, R., Goldman, N., Pedersen, A.M., 2000. Codon-substitution models for heterogeneous selection pressure at amino acid sites. *Genetics* 155, 431–449.
- Yang, W., Cheng, T., Ye, M., Deng, X., Yi, H., Huang, Y., Tan, X., Han, D., Wang, B., Xiang, Z., Cao, Y., Xia, Q., 2011. Functional divergence among silkworm antimicrobial peptide paralogs by the activities of recombinant proteins and the induced expression profiles. *PLoS One* 6, e18109.

mature peptide

B. lucorum	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAACAAATATTTAAAA
B. t. xanthopus	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAACAAATATTTAAAA
B. terrestris	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAACAAATATTTAAAA
B. lapidarius	TGGCCATATCCATTACCAAAC	GTACGTATATCTTAACATTTGAACAATAAATATTTAAAA
B. bohemicus	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAATAAATATTTAAAA
B. maxillosus	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAATAGATATTTAAAA
B. perezi	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAATAAATATTTAAAA
B. vestalis	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTAAACAATAAATATTTAAAA
B. rupestris	TGGCCATACCCATTACCAAAC	GTACGTATATTTTAACATTTAAACAATAAATATTTAAAA
B. campestris	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACGTTTAAACAATAAATATTTAAAA
B. ruderus	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATTTGAACAACAATAAATATTTAAAA
B. pascuorum	TGGCCATACCCATTACCAAAC	GTACGTATATCTTAACATCTAAACAATAAATATTTAAAA

***** .***** .***** .***** .***** .***** .*****

B. lucorum	GTATTTACTAGACTAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. t. xanthopus	GTATTTACTAGACTAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. terrestris	GTATTTACTAGACTAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. lapidarius	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTGTTATGAAGT-ATATTCAAGAGT
B. bohemicus	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. maxillosus	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. perezi	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. vestalis	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. rupestris	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT
B. campestris	GTATTTACTAGACAGAACTTTATTTAAAAGGAATATTCTTGTGAAGTAAATTTCAAGAGT
B. ruderus	GTATTTACTAGACTGAAACTTTATTTAAAAGGAATATTGTTATAAAGT-ATATTCAAGAGT
B. pascuorum	GTATTTACTAAACTGAAACTTTATTTGAAAGGAATATTCTTATGAAGT-ATATTCAAGAGT

***** .**:.***** .***** .***** .***** .***** .*****

Intron 1

B. lucorum	TATATGTATAAAGTCCCTGAGAATACACACTTTCATAGAATCGGAATATAGATATAGGTT
B. t. xanthopus	TATATGTATAAAGTCTCTGAGAATACACACTTTCATAGAATCGGAATATAGATATAGGTT
B. terrestris	TATATGTATAAAGTCTCTGAGAATACACACTTTCATAGAATCGGAATATAGATATAGGTT
B. lapidarius	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAAATCGGAATACAGATATAGGTT
B. bohemicus	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAAATCGGAATATAGATATAGGTT
B. maxillosus	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAAATCGGAATATAGATATAGGTT
B. perezi	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAAATCGGAATATAGATATAGGTT
B. vestalis	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAAATCGGAATATAGATATAGGTT
B. rupestris	TA----TATAAAGTCCCTGAGAATACACAATTTTATAAATCGGAATATAAATATAAGTT
B. campestris	TA----TATAAAGTCCCTGAAAATACACAATTTTATAGAATTTGGAATATAGATATAGGTT
B. ruderus	TA----TATAAAGTCCCTGAGTATACACAATTTTATAAAATCGGAATATAGA-----
B. pascuorum	TA----TATAAAGTCCCTGAAAATACACAATTTTATAAAATCGGAATATAGATATAGGTT

** ***** .***** .***** .***** .***** .***** .*****

B. lucorum	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. t. xanthopus	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. terrestris	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. lapidarius	GTTTATGTAGTTATAAAT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. bohemicus	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. maxillosus	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. perezi	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. vestalis	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. rupestris	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. campestris	GTTTATGTAGTATATAAATT-TTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG
B. ruderus	-----TAGTATATAAATCTTTTTTACTAAAAATTTAAAAATTTCTCAAAATGTTAATG
B. pascuorum	GTTTATGTAGTATATAAAGTTTTTTTTTATTAATAAATTTCAAAATTTCTCAAAATGTTAATG

****.***** *****:***** .***** .***** .***** .*****

B. lucorum	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. t. xanthopus	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. terrestris	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. lapidarius	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. bohemicus	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. maxillosus	CATGACTTTGAAATTAAGAATGCTTAACATTATATTTATA--TAAATTAGAAAACGTGCGG
B. perezi	CATGACTTTGAAATTAAGAATGCTTAACATTATATTTATA--TAAATTAGAAAACGTGCGG
B. vestalis	CATGACTTTGAAATTAAGAATGCTTAACATTATATTTATA--TAAATTAGAAAACGTGCGG
B. rupestris	CATGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGCGG
B. campestris	CACGACTTTGAAATTAAGAATGCTTAACATTATGTTTATATGTAAATTAGAAAACGTGCGG
B. ruderus	CATGACTTTGAAATTAAGAATGCTTAACACCATATTTATA--TAAATTAGAAAACGTGCGG
B. pascuorum	CACGACTTTGAAATTAAGAATGCTTAACATCATATTTATA--TAAATTAGAAAACGTGGG

** ***** .***** .***** .***** .***** .***** .*****

B. lucorum	GGCGAATATA-CTTTTTACAGAATATTTCCAGTAATATTCAATTTTCATCTAAATAGAAAA
B. t. xanthopus	GGCGAATATA-CTTTTTACAGAATATTTCCAGTAATATTCAATTTTCATCTAAATAGAAAA
B. terrestris	GGCGAATATA-CTTTTTACAGAATATTTCCAGTAATATTCAATTTTCATCTAAATAGAAAA
B. lapidarius	GGCGAATATA-CTTTTTACAGAATATTTCCAGTAATATTCAATTTTCATCTAAATAGAAAA
B. bohemicus	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. maxillosus	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. perezi	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. vestalis	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. rupestris	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. campestris	AGCGAATATA-TTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTGAATAGAAAA
B. ruderus	GACGAATATATTTTTTACAGAATATTTCCAATAATATTCAATTTTCATCTAAATAGAAAA
B. pascuorum	AGCGAATA--CTTTTTACAGAATATTTCCAGTAATATTCAATTTTATCTAAATAGAGAA

..***** ***** .***** .***** .***** .***** .*****

B. lucorum	TTTTAAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAAACA
B. t. xanthopus	TCTTAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAAACA
B. terrestris	TCTTAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAAACA
B. lapidarius	TCTTAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAGATA
B. bohemicus	TCTTAAAAATCTTAAGAACACTATTAACGTTTCGTAAAAATTATTACGAAAAATATTTAGACA
B. maxillosus	TCTTAAACAATCTTAAGAACACTATTAACGTTTCGTAAAAATTATTACGAAAAATATTTAGACA
B. perezi	TCTTAAACAATCTTAAGAACACTATTAACGTTTCGTAAAAATTATTACGAAAAATATTTAGACA
B. vestalis	TCTTAAACAATCTTAAGAACACTATTAACGTTTCGTAAAAATTATTACGAAAAATATTTAGACA
B. rupestris	TCTTAAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTACTACGAAAAATATTTAGACA
B. campestris	TCTTAAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAGACA
B. ruderalis	TCTTAAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAGACA
B. pascuorum	TCTTAAACAATCTTAAGAACACTATTAACGTTTGTAAAAATTATTACGAAAAATATTTAGACA
	* **** * :***** * * ***** ***** *
	mature peptide
B. lucorum	TTATCGGATATTATATATATATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. t. xanthopus	TTATCGGATATTATATATATATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. terrestris	TTATCGGATATTATATATATATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. lapidarius	TTATCGGATATT--ATATATATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. bohemicus	TTATCGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. maxillosus	TTATCGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. perezi	TTATCGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. vestalis	TTATCGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. rupestris	TTATCGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. campestris	TTATTGGATATTAAATATATATAATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
B. ruderalis	TTATCGGATATTATATATATATTTGTTTTATTACAGCTGGCCACTAATCACAACAAA
B. pascuorum	TTATCGGATATTATATATATATTTGTTTTATTACAGCTGGTCATTAATCACAACAAA
	**** * :***** * * ***** ***** *
B. lucorum	TCAAGAAACGTTCAAACAGTTAAACTCGACAGATCATCAGAGATTATATCTACAATGTAC
B. t. xanthopus	TCAAGAAACGTTCAAACAGTTGAACCTCGACAGATCATCAGAGATTATATCTACAATGTAC
B. terrestris	TCAAGAAACGTTCAAACAGTTGAACCTCGACAGATCATCAGAGATTATATCTACAATGTAC
B. lapidarius	TCAAGAAACGTTCAAACAGTTGAACCTCGACAGATCATCAGAGATTATATCTACAATGTAC
B. bohemicus	TCAAGAAACGTTGAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. maxillosus	TCAAGAAACGTTGAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. perezi	TCAAGAAACGTTGAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. vestalis	TCAAGAAACGTTGAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. rupestris	TCAAGAAACGTTCAAACAATTCACCTTGACAAATCATCAGAGATTATATCTATAATGTAG
B. campestris	TCAAGAAACGTTCAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. ruderalis	TCAAGAAACGTTCAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAG
B. pascuorum	TCAAGAAACGTTCAAACAATTCACCTTGACAAATCATCAGAGATTATATCTACAATGTAT
	***** * :***** * * ***** ***** *
B. lucorum	CAACAAATTATTATAAATTATAAATTGTTATTAAGTTATACACTGATTAAAAATGATCCTGA
B. t. xanthopus	CAACAAATTATTATAAATTATAAATTGTTATTAAGTTATACACTGATTAAAAATGATCCTGA
B. terrestris	CAACAAATTATTATAAATTATAAATTGTTATTAAGTTATACACTGATTAAAAATGATCCTGA
B. lapidarius	CGACGAATTATTATAAATTATAAATTG--AATTATGTTATACACTGATTAAAAATGATCCTGA
B. bohemicus	CGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACACTCATTACATTGATCCTGA
B. maxillosus	CGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACACTCATTACATTGATCCTGA
B. perezi	CGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACACTCATTACATTGATCCTGA
B. vestalis	CGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACACTCATTACATTGATCCTGA
B. rupestris	CGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACACTCATTACATTGATCCTGA
B. campestris	TGACAAATTATCATAAATATAAATTA--AATTAAAGTTATACATTACATTGATCCTGA
B. ruderalis	CGATGAATTATTATAAATATAAATTG--AATTAAAGTTATACACTGATTAAAAATGATCCTGA
B. pascuorum	CGACGAATTATTATAAATATAAATG--AATTAAAGTTATACGCTCATTAAAAATGATCCTGA
	* :***** * * ***** ***** *
B. lucorum	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. t. xanthopus	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. terrestris	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. lapidarius	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. bohemicus	TTTATCTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. maxillosus	TTTATCTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. perezi	TTTATCTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. vestalis	TTTATCTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. rupestris	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTTATTTA
B. campestris	TTTATTTTGAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. ruderalis	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
B. pascuorum	TTTATTTTCAATAAAATAGAAACTATCCTGAAACATAGTGTAAATTAATGTTCAATTA
	***** * :***** * * ***** ***** *
B. lucorum	ACACTAAATTGATCACTGATATTTCCCTTACGAA--ATAATCGAAGTTAATGAAGTACGTGC
B. t. xanthopus	ACACTAAATTGATCACTGATATTTCCCTTACGAA--ATAATCGAAGTTAATGAAGTACGTGC
B. terrestris	ACACTAAATTGATCACTGATATTTCCCTTACGAA--ATAATCGAAGTTAATGAAGTACGTGC
B. lapidarius	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. bohemicus	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. maxillosus	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. perezi	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. vestalis	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. rupestris	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. campestris	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. ruderalis	ACACTAAATCGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC
B. pascuorum	ACACTAAATTGATCACTGATATTTCCCTTACGAACATAATCGAAGTTAATGAAGTACGTGC

```

*****  *****  *****  *****  *****  *****
B.lucorum      GCGCATCTTTCTTTTCGTCAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.t.xanthopus  GCGCATCTTTCTTTTCGTCAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.terrestris   GCGCATCTTTCTTTTCGTCAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.lapidarius   GCGCATCTTTCTTTTCGTCAGATCATGTTTAACATTATCAAATGGAGAGGTACATAGTA
B.bohemicus    GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.maxilloso    GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.perezi       GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.vestalis     GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.rupestris    GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.campestris   GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.ruderatus    GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
B.pascuorum    GCGTATCTTTCTTTTCGTTAGATCATGTTTAACATTATCAAATGGAGAGGTACAAAGTA
** . *****  *****  *****  *****  *****  *****

B.lucorum      TTTTACCTTGGATTGCGCTTTAATAGAAAACCTGTGTTTATTGTTAAATAAAAGTAAT
B.t.xanthopus  TTTTACCTTGGATTGCGCTTTAATAGAAAACCTGTGTTTATTGTTAAATAAAAGTAAT
B.terrestris   TTTTACCTTGGATTGCGCTTTAATAGAAAACCTGTGTTTATTGTTAAATAAAAGTAAT
B.lapidarius   TTTTACCTTGGATTGCGCTTTAATAGAGAACTCGTGTTTATTCTTAAATAAAAGTAAC
B.bohemicus    TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCGTATTATTCTTAAATAAAAGTAAT
B.maxilloso    TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCGTATTATTCTTAAATAAAAGTAAT
B.perezi       TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCATATTATTCTTAAATAAAAGTAAT
B.vestalis     TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCATATTATTCTTAAATAAAAGTAAT
B.rupestris    TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCGTATTATTCTTAAATAAAAGTAAT
B.campestris   TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCGTATTATTCTTAAATAAAAGTAAT
B.ruderatus    TTTGTACCTTAGATTGCGCTTTAATAGAGAACTCGTGTTTATTCTTAAATAAAAGTAAT
B.pascuorum    TTTGTATCTTAGGTTT-----
*** ** ** *.***

B.lucorum      AATAGAAAATAGAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.t.xanthopus  AATAGAAAATAGAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.terrestris   AATAGAAAATAGAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.lapidarius   AGTAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.bohemicus    AATAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATATCATTTTA
B.maxilloso    AATAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATATATTTTA
B.perezi       AATAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATATATTTTA
B.vestalis     AATAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATATATTTTA
B.rupestris    AATAGAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.campestris   AATAAAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTTA
B.ruderatus    AATAGAAAATAGAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATA--TCATTTTA
B.pascuorum    ----GAAAATATAAAATATTCGTAGCTTTTAATATTCTTCTCCCAAAATATGTCATTTT-
. *****  *****  *****  *****  *****  *****

B.lucorum      ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAC
B.t.xanthopus  ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAC
B.terrestris   ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAC
B.lapidarius   ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAC
B.bohemicus    ATATAAGAAAGAAACGCATAAAATACACCAGTTTATTTTCATTAGAAAATTACGATATAC
B.maxilloso    ATATAAATAAGAAACGCATAAAATACACCAGTTTATTTTCATTAGAAAATTACGATATAC
B.perezi       ATATAAATAAGAAACGCATAAAATACACCAGTTTATTTTCATTAGAAAATTACGATATAC
B.vestalis     ATATAAATAAGAAACGCATAAAATACACCAGTTTATTTTCATTAGAAAATTACGATATAC
B.rupestris    ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAG
B.campestris   ATATAAGAAAGAAACGCATAAAATACAGCAGTTTATTTTCATTGAAAATTACGATATAC
B.ruderatus    ATATAAGGAAGAAACGCATAAAATACAGCAGTTTATTTTATTGAAAATTACGATATAC
B.pascuorum    -----CCAAATACAGCAGTTTATTTTCATTGAAAATTACGATGTAC
. *****  *****  *****  *****  *****  *****

B.lucorum      ATAGGTATGGGATGATATCGATCTGCTGGTACAAACGAAAAGGG
B.t.xanthopus  ATAGGTATGGGATGATATCGATCTGCTGGTACAAACGAAAAGGG
B.terrestris   ATAGGTATGGGA-----
B.lapidarius   ATAGGTATGGGATGATATCGATCTGCTGGTACAAACGAAA-----
B.bohemicus    GTAGGTATGGGATGATATCGATCTAGTGGTACAAACGAAAAGAG
B.maxilloso    ATAGGTATGGGATAATATCGATCTAGTGGTACAAACGAAAAGAG
B.perezi       ATAGGTATGGGATAATATCGATCTAGTGGTACAAACGAAAAGAG
B.vestalis     ATAGGTATGGGATAATATCGATCTAGTGGTACAAACGAAAAGAG
B.rupestris    ATAGGTATGGGATGATATCGATCTAGTGATACAAACGAAAAGAG
B.campestris   ATAGGTATGGGATGATATCGATCTAG--GGTACAAACGAAAAGAG
B.ruderatus    ATAGGTATGGGGTATATCGATCTGCTGGTACAAACGAAAAGGG
B.pascuorum    ATAGGTATGGGATGATATCGATCTGCTGGTACAAACGAAAAGGG
. *****

```

Figure S1: *Abaecin* sequence alignment using MAFFT version 6 (Katoh and Toh, 2010; Katoh et al., 2002) and annotated according to Choi et al., (2008).

B. lucorum	AGCTGCTGTTTTAATGCTCCACCGTCTTCGCCTTTCTATTGAGAAATATCCGAGCGTTAA
B. terrestris	-----ATATCCGAGCGTTAA
B. t. xanthopus	AGCTGCTGTTTTAATGCTCCACCGTCTTCGCCTTTCTATTGAGAAATATCCGAGCGTTAA
B. pascuorum	AGCTGCTGTTTTAATGCTCCATCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA
B. ruderus	AGCTGCTGTTTTAATGCTCCATCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA
B. bohemicus	AGCTGCTGTTTTAATGCTCCACCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA
B. vestalis	AGCTGCTGTTTTAATGCTCCACCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA
B. campestris	-----
B. rupestris	AGCTGCTGTTTTAATGCTTCACCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA
B. lapidarius	AGCTACTGTTTTAATGCTCCGTCATCTTCGTCTTTCTATTGAGAAATATCCTAGCGTTAA
B. maxillosus	AGCTGCTGTTTTAATGCTCCACCATCTTCGTCTTTCTATTGAGAAATATCCAAGCGTTAA
B. perezi	AGCTGCTGTTTTAATGCTCCACCATCTTCGTCTTTCTATTGAGAAATATCCGAGCGTTAA

B. lucorum	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. terrestris	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. t. xanthopus	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. pascuorum	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. ruderus	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. bohemicus	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGTAGCAGACAGAAGGCTT
B. vestalis	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGTAGCAGACAGAAGGCTT
B. campestris	-----
B. rupestris	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGTAGCAGACAGAAGGCTT
B. lapidarius	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGCAGCAGACAGAAGGCTT
B. maxillosus	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGTAGCAGACAGAAGGCTT
B. perezi	TTCTCGATTCCTTTGCCAATTACCGTGTTTCATCTATTATACGTAGCAGACAGAAGGCTT

pro peptide

B. lucorum	AAAGACGGTATTTAAATTTTCAGAGGAGGAGTACGAGCTCCTCGAACAAAGCTGGTATCGAG
B. terrestris	AAAGACGGTATTTAAATTTTCAGAGGAGGAGTACGAGCTCCTCGAACAAAGCTGGTATCGAG
B. t. xanthopus	AAAGACGGTATTTAAATTTTCAGAGGAGGAGTACGAGCTCCTCGAACAAAGCTGGTATCGAG
B. pascuorum	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGGTATCGAG
B. ruderus	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGATATCGAG
B. bohemicus	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAG
B. vestalis	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAG
B. campestris	-----AGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAA
B. rupestris	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAG
B. lapidarius	AAAAACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGATATCGAG
B. maxillosus	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAG
B. perezi	AAAGACGGTATTTAAATTTTCAGAGGAGGAATACGAGCCCTCGAACACGCTGCTATCGAG

* * * * * , * * * * * * * * * * , * * * * * , * * * * * ,

B. lucorum	GAACGTGCCGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCA
B. terrestris	GAACGTGCCGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCA
B. t. xanthopus	GAACGTGCCGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCA
B. pascuorum	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. ruderus	GAACGTGCTGTTAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. bohemicus	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. vestalis	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. campestris	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. rupestris	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. lapidarius	GAACGTGCTGACAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. maxillosus	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT
B. perezi	GAACGTGCTGATAGACAAAGAAAGAGTGACCTGCGACCTTCTCTCCATCAAAGGAGTCGCT

* * * * * * : * * * * * , * :

B. lucorum	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. terrestris	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. t. xanthopus	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. pascuorum	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. ruderus	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. bohemicus	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. vestalis	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. campestris	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. rupestris	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. lapidarius	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. maxillosus	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG
B. perezi	GAACATAGTGCTTGCGCTGCCAACTGTCTCAGCATGGGCAAAGCTGGAGGTCGCTGCGAG

* :

| | |
|-----------------|--|
| B. lucorum | AACGGAGTCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. terrestris | AACGGAGTCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. t. xanthopus | AACGGAGTCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. pascuorum | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. ruderus | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. bohemicus | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. vestalis | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. campestris | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. rupestris | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. lapidarius | AACGGAGTCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. maxillosus | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |
| B. perezi | AACGGAATCTGCCTTTGTGCGAAGTACGTGATTCCCTTTAATTTTCATCTGCCCAAGTTGTT |

* * * * * , *

mature peptide

B. lucorum TGAACCTAAACGACCAATTTTGAATGTTTTGCCGCGACTATTTCGACGCTCTGTTGATCGAA
B. terrestris TGAACCTAAACGACCAATTTTGAATGTTTTGCCGCGACTATTTCGACGCTCTGTTGATCGAA
B. t. xanthopus TGAACCTAAACGACCAATTTTGAATGTTTTGCCGCGACTATTTCGACGCTCTGTTGATCGAA
B. pascuorum TGAACCTAAACAACCAATCTTGAACGTTTTGCT-----ACGCTGTTGATCGAA
B. ruderatus CGAACTGAAACAACCAATCTTCAATGTTTTGCT-----ACGCTGTTGATCGAA
B. bohemicus TGAACCTAAACAACCAATCTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
B. vestalis TGAACCTAAACAACCAATCTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
B. campestris TGAACCTAAACAACCAATCTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
B. rupestris TGAACCTAAACAACCAATCTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
B. lapidarius TGAACCTAAACGACCAATCTCGAATGTTTTGCTACGACTATTTCGACGCTCTGTTGATCGAA
B. maxillosus TGAACCTAAACAACCAATTTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
B. perezi TGAACCTAAACAACCAATTTTGAATGTTTTGCT-----ACGCTGTTGGTCGAA
***** **

B.lucorum GGAAAATTCTCGCAGGTTAAACGATTATTAAATATAGCCAACCTTCAGTTTATTAAATTT
 B.terrestris GGAAAATTCTCGCAGGTTAAACGATTATTAAATATAGCCAACCTTCAGTTTATTAAATTT
 B.t.xanthopus GGAAAATTCTCGCAGGTTAAACGATTATTAAATATAGCCAACCTTCAGTTTATTAAATTT
 B.pascuorum GGAAAATTCTCGCAGGTTA---GATTATTAAATATAGCTAAGTTTCAGTTTATTAAATTT
 B.ruderatus GGAAAATTCTCGCAGGTTAGACGATTATTAAATATAGCCAAGTTTCAGTTTATTAAATTT
 B.bohemicus GGAAAATTCTCGCAGGTTAGACGATTATTAAATATAGCCAAGTTTCGGTTTATTAAATTT
 B.vestalis GGAAAATTCTCGCAGGTTAGACGATTATTAAATATAGCCAAGTTTCGGTTTATTAAATTT
 B.campestris AGAAAATTCTCGCAGGTTAGACGATTATTAAATATAG-----
 B.rupestris GGAAAATTCTCGCAGGTTAGACGATTATTAAATATAGCCAAGTTTCGGTTTATTAAATTT
 B.lapidarius GGAAAATTCTCGCAGGTTAAACGATTATTAAATACAGCCAAGTTTCAGTTTGTAAATGT
 B.maxillosus -----TTATTAAATTT
 B.perezi -----

B.lucorum CCAACGAGTTGAATTTGAACGACGATTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.terrestris CCAACGAGTTGAATTTGAACGACGATTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.t.xanthopus CCAACGAGTTGAATTTGAACGACGATTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.pascuorum CGAACGAAATTGAATTTGAACGACGACTAGGTTTTTCAGTACTGTAGCGTAGATACAATAAT
 B.ruderatus CGAACGAGTTGAATTTGAACGACGACTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.bohemicus CGAACGAGTTGAATTTGAACGACGACTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.vestalis CGAACGAGTTGAATTTGAACGACGACTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.campestris -----
 B.rupestris CGAACGAGTTGAATTTGAACGACGACTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.lapidarius CGAACGAGTTGAATTTGAACGACGATTAGGTTTTTCGGTATTATAGCGTAGATCCAATAAT
 B.maxillosus CGAACGAGTTGAATTTGAACGACGACTAGGTTTTTCAGTACTATAGCGTAGATTCAATAAT
 B.perezi -----

B.lucorum CTTTTGGTAAATTCAATGCCCATTTGAAAATTTTCGTCCTACGAAATGTAACCTTTGTTTT
 B.terrestris CTTTTGGTAAATTCAATGCCCATTTGAAAATTTTCGTCCTACGAAATGTAACCTTTGTTTT
 B.t.xanthopus CTTTTGGTAAATTCAATGCCCATTTGAAAATTTTCGTCCTACGAAATGTAACCTTTGTTTT
 B.pascuorum CTTTTAGTAAATTCAATGCCCGTTGAAAATTTTCGTGATACGAAATGTAACCTTTG-TTT
 B.ruderatus CTTTCAGTAAATTCAATGCTCGTTGAAAATTTTCGTGATACGAAACGTAACCTTTGTTTT
 B.bohemicus CTTTTAGTAAATTAAATGCCCACTGAAAATTTTCGTGCATACGAAATGTAACCTTTG-TTT
 B.vestalis CTTTTAGTAAATTAAATGCCCACTGAAAATTTTCGTTATACGAAATGTAACCTTTG-TTT
 B.campestris -----
 B.rupestris CTTTTAGTAAATTCAATGCCCGCTGAAAATTTTCGTGATACGAAATGTAACCTTTG-TTT
 B.lapidarius CTTTTAGTAAATTCAATGCCCGTTGAAAATTTTCGTGATACGAAATGTAACCTTTGTTTT
 B.maxillosus CTTTTAGTAAATTAAATGCCCACTGAAAATTTTCGTTATACGAAATGTAACCTTTG-TTT
 B.perezi -----

B.lucorum TTATAGTACCGCAGACGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.terrestris TTATAGTACCGCAGACGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.t.xanthopus TTATAGTACCGCAGACGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.pascuorum TTATAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAGTTCGCTTGGAAGTTCT
 B.ruderatus TTATAGTACCACAGACGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.bohemicus TTATAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.vestalis TTATAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.campestris -----
 B.rupestris TTATAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.lapidarius TTAAAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.maxillosus TTATAGTACCACAGATGTTTCAGGAGAAAATTCGTGTTTTAATTCGCTTGGAAGTTTCT
 B.perezi -----CGTGTTTTAATTAGCTTGGAAGTTTCT

B.lucorum ATCGTTCGTGTTGGGCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTACCAT
 B.terrestris ATCGTTCGTGTTGGGCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTACCAT
 B.t.xanthopus ATCGTTCGTGTTGGGCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTACCAT
 B.pascuorum ATTGTTTCGTGTTGGATCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTACCAT
 B.ruderatus ATTGTTTCGTGTTGGATCTCGTGTCACGTTGATTATCAACAGCTCAGCTGTTATCAT
 B.bohemicus ATTGTTTCGTGTTGAATCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTATCAT
 B.vestalis ATTGTTTCGTGTTGAATCTCGTGTCACGTTGATTATCGATAGCTCAGCTCTTATCAT
 B.campestris -----
 B.rupestris ATTGTTTCGTGTTGAATCTCGTGTCACGTTGATTATCGATATCTCAGCTGTTATCAT
 B.lapidarius ATTGTTTCGTGTTGGACCTCGTGTCACGTTGATTATCGATAGCTCAGCTGTTACCAT
 B.maxillosus ATTGTTTCGTGTTGAATCTCGTGTCACGTTGATTATCGATAGCTCAGCTCTTATCAT
 B.perezi ATTGTTTCGTGTTGAATCTCGTGTCACGTTGATTATCGATAGCTCAGCTCTTATCAT

B.lucorum AGATGAAGTTCTACTACATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.terrestris AGATGAAGTTCTACTACATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.t.xanthopus AGATGAAGTTCTACTATATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.pascuorum AGATGAAGTTCTACTAGATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.ruderatus AGATGAAGTTCTACTGGATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.bohemicus AGATGAAGTTTTTACTAAATGAAAATTCATAAGTTTATTACTAAAACGAGACGAAAGCA
 B.vestalis AGATGAAGTTCTTACTAGATGAAAATTCATAAGATTTATTACTAAAACGAGACGAAAGCA
 B.campestris -----
 B.rupestris AGATGAAGTTCTTACTAGATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGTA
 B.lapidarius AGACGAAGTTCTATTAGATGAAAATTCATAAGATTTGTTACTAAAACGAGGCGAAAGCA
 B.maxillosus AGATGAAGTTCTTACTAGATGAAAATTCATAAGATTTATTACTAAAACGAGACGAAAGCA

| | |
|---------------|---|
| B.perezi | AGATGAAGTTCTTACTAGATGAAAATTCATAAGATTTATTACTAAAACGAGACGAAAGCA |
| | |
| B.lucorum | AAATTTAGAGAATCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCGAAGATA |
| B.terrestris | AAATTTAGAGAATCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCGAAGATA |
| B.t.xanthopus | AAATTTAGAGAATCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCGAAGATA |
| B.pascuorum | AAGTTTAGAGAGTCAGGAAGCTGATTATTGTTGCTGTGCGTAATCAAATACCCAAAGATA |
| B.ruderatus | AAATTTAGAGAGTCAGGAAGCTGATTATTATTGTTGCTGTGCGTAATCGAATATCCAAAGATA |
| B.bohemicus | AAATTTAGAGAGTCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATACCCAAAGATA |
| B.vestalis | AAATTTAGAGAGTCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCAAAGATA |
| B.campestris | ----- |
| B.rupestris | AAATTTAGAGAGTCAGGAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCAAAGATA |
| B.lapidarius | AAGTTTAGAGAGTCAGGAAGCTGATTATTATTGCTGTGCGTAATCAAATATCCAAAGATA |
| B.maxilloso | AAATTTAGAGAGTCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCAAAGATA |
| B.perezi | AAATTTAGAGAGTCAGCAAGCTGATTATTGTTGCTGTGCGTAATCAAATATCCAAAGATA |
| | |
| B.lucorum | ATCTCGCGTCATAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATTGGACG |
| B.terrestris | ATCTCGCGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATTGGACG |
| B.t.xanthopus | ATCTCGCGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATTGGACG |
| B.pascuorum | ATCTCGTGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAA-----CTGGACG |
| B.ruderatus | ATCTCGTGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTTACCTGGACG |
| B.bohemicus | ATCTCGTGTCTCAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATTGGACG |
| B.vestalis | ATCTCGTGTCTCAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATCTGGACG |
| B.campestris | ----- |
| B.rupestris | ATCTCGTGTCTCAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATTGGACG |
| B.lapidarius | ATCTCGTGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATCTGGACG |
| B.maxilloso | ATCTCGTGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATCTGGACG |
| B.perezi | ATCTCGTGTCTTAATTTTGAAGTATTGTTGGAGGTAGCTAACTTTCGTGTCATCTGGACG |
| | |
| B.lucorum | TGTTGTAGATATTTCAAATGTT |
| B.terrestris | TGTTGTAGATATTTCAAATGTT |
| B.t.xanthopus | TGTTGTAGATATTTCAAATGTT |
| B.pascuorum | TGTTATAGATATTTCAAATGTT |
| B.ruderatus | TGTTACAGATATTTGAAATGTT |
| B.bohemicus | TGTTATAGATATTTCATATGTT |
| B.vestalis | TGTTATAGATATTTCATATGTT |
| B.campestris | ----- |
| B.rupestris | TGTTATAGATATTTCAAATGTT |
| B.lapidarius | TGTTATAGATACTTCAAATGTT |
| B.maxilloso | TGTTATAGATATTTCATATGTT |
| B.perezi | TGTTATAGATATTTCATATGTT |

Figure S2: *Defensin-1* sequence alignment using MAFFT version 6 (Katoh and Toh, 2010; Katoh et al., 2002) and annotated according to Choi et al., (2008)

B.lucorum GTAAGAACTTTACTATATAATTAATTTTAAAAATTGCAACTTAAAAATTAGCACTAAATGT
 B.terrestris GTAAGAACTTTACTATTTAATTAATTTGTAATAATTGCTACTTAAAAATTAGCACTAAATGT
 B.t.xanthopus GTAAGAACTTTACTATTTAATTAATTTGTAATAATTGCTACTTAAAAATTAGCACTAAATGT
 B.pascuorum GTAAGAACTTTACTATATAATTAATTTTAAAAATTGCTACTTAAAAATTAGCACTAAATGT
 B.ruderatus GTAAGAAATTCACATATAATTAATTTTAAAAATTGCTACCTAAAAATTAGCACTAAATGC
 B.bohemicus GTAAGAACTTTACTATACAATTAATTTTAAAAATTGCTACTTAAATATTAGCATTAAATGT
 B.maxilloso B.maxilloso B.maxilloso B.maxilloso B.maxilloso B.maxilloso B.maxilloso
 B.perezi GTAAGAACTTTACTATACAATTAATTAATTAATTTGCTACTTAAATATTAGCATTAAATGT
 B.vestalis GTAAGAACTTTACTATACAATTAATTAATTAATTTGCTACTTAAATATTAGCATTAAATGT
 B.rupestris GTAAGAACTTTACTATACAATTAACCTTAAAAATTGCTACTTAAATATTAGCATTAAATGT
 B.lapidarius GTAAGAACTTTACTATATACTTAATTTTAAAAATTGCTACTTAAAAATTAGCACCAAATGT
 B.campestris GTAAGAAATTTACTATACAATTAATTTTAAAAATTGCTACTTAAATATTAGCATTAAATGT
 ***** ** *****: *.**** : *.:*****.* *****:***** *****

B.lucorum GATTGAAACAATATTACAATAGGATA-CTAGTTGGATAATTAGTACTTTTCATCCAATAGA
 B.terrestris GATTGAAACAATATTACAATAGGATA-CTAGTTGGATAATTAGTACTTTTCATCCAATAGA
 B.t.xanthopus GATTGAAACAATATTACAATAGGATA-CTAGTTGGATAATTAGTACTTTTCATCCAATAGA
 B.pascuorum GATTGAAATAATATTACAATAGAATA-CTAGTTGGATAATTA-----
 B.ruderatus GATTGAAATAATATTACGATGGAATA-CTAGTTGGATAATTAGTACTTTTCATTTAATAAA
 B.bohemicus GACTGAAATAATATGACAATGGAATA-GTCTTTAGATAATTAGTACTTTTCATCTAATAAA
 B.maxilloso GACTGAAATAATATGACAATGGAATA-GTCTTTAGATAATTAGTACTTTTCATCTAATAAA
 B.perezi GACTGAAATAATATGACAATGGAATA-GTCTTTAGATAATTAGTACTTTTCATCTAATAAA
 B.vestalis GATTGAAATAATATGACAATGGAATA-GTCTTTAGATAATTAGTACTTTTCATCTAATAAA
 B.rupestris GATTGAAATAATATGACAATGGAATA-GTATTTAGATAATTAGTACTTTTCATCTAATAAA
 B.lapidarius GATTGAAATAATATGACAATGGAATA-GTATTTGGATAATTAGTACTTTTCACCTAATAAA
 B.campestris GACTGAAATAATATGACAATGGAATATATATTAGATAATTAGTACTTTTCATCTAATAAA
 ** ***** ***** *.**.*.*** *.**.*.*****

B.lucorum TGGTAATGTCTAAATCTGTTACTCTGCAATAATTTGAAAATTGCGTATGGATTGGTTACA
 B.terrestris TGGTAATGTTTTAAATCTGTTACTCTGCAATAATTTGAAAATTGCTTATGGATTGGTTACA
 B.t.xanthopus TGGTAATGTTTTAAATCTGTTACTCTGCAATAATTTGAAAATTGCTTATGGATTGGTTACA
 B.pascuorum --GTAATTTTTTAAATCTGTAACCTCGGCAATAATTTGAAAATTGCAATGATTGGTTACA
 B.ruderatus TGGTAATTTTTTAAATCTGTAACCTCTGCAATAATTTGAAAATTGCGTATGGATCGGTTACA
 B.bohemicus TGGTAATTTTTGTATCTGTAACCTTTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 B.maxilloso TGGTAATTTTTGTATCTGTAACCTTTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 B.perezi TGGTAATTTTTGTATCTGTAACCTTTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 B.vestalis TGGTAACTTTTGTATCTGTAACCTTTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 B.rupestris TGGTAATTTTTGTATCTGTAACCTGTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 B.lapidarius TAGTAATTTTTTAAATCTGTAACCTCTGCAATAATTTGAAAATTGCGTATGGATT-----
 B.campestris TGGTAATTTTTGTACCTGTAACCTCTGCAATAATTTGAAAATTGCGTATGTAGTGGTTACA
 **** * *.:* *****.* *****.* ***** *

B.lucorum AATATACAAATTAAAAATGTCATCAAGCGAGAATTGTTTACGAAAATCTATAATAGAACG
 B.terrestris AATATACAAATTAAAAATGTCATCAAGCGAGAATTGTTTACGAAAATCTATAATAGAACG
 B.t.xanthopus AATATACAAATTAAAAATGTCATCAAGCGAGAATTGTTTACGAAAATCTATAATAGAACG
 B.pascuorum AATATACAACCTTAAAAATATCATTAAGCGAGAATTGTTTAGGAAAATCTATAATAGAACG
 B.ruderatus AATATACAACCTTAAAAATTTTATTAAAGCGAAAATTGTTTACGAAAATTTATAATAGAACG
 B.bohemicus AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGTTTATAAAAAATCTATAATAGAACG
 B.maxilloso AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGTTTATGAAAATCTATAATAGAACG
 B.perezi AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGTTTATAAAAAATCTATAATAGAACG
 B.vestalis AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGTTTATAAAAAATCTATAATAGAACG
 B.rupestris AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGTTTATAAAAAATCTATAATAGAACG
 B.lapidarius -----
 B.campestris AATATACAACCTTAAAAATGTCATTAAGCAAGAATTGATTATGAAAATCTAGAATAGAGCG

B.lucorum AATTTGAGCATTTTGATGAATTTTCCAACGCTACATCTTAACTTAACATTTTTATCCTT
 B.terrestris AATTTGAGCATTTTGATGAATTTTCCAACGCTACATCTTAACTTAACATTTTTATCCTT
 B.t.xanthopus AATTTGAGCATTTTGATGAATTTTCCAACGCTACATCTTAACTTAACATTTTTATCCTT
 B.pascuorum AATTTGAGAATTTTCAAGAATTTTCCAACGCTACATCTTAACTTAACATTTTTATTCTT
 B.ruderatus AATTTGAGAATTTTGAAGAATTTTCCAACGCTGATCTTAACTTAACATTTTTATTCTT
 B.bohemicus AATTTGAGAATTTTCAAGAATTTTCCAATGCTACATCTTAACTTAACATTTTTATTCTT
 B.maxilloso AATTTGAGAATTTTCAAGAATTTTCCAATGCTACATCTTAACTTAACATTTTTATTCTT
 B.perezi AATTTGAGAATTTTCAAGAATTTTCCAATGCTACATCTTAACTTAACATTTTTATTCTT
 B.vestalis AATTTGAGAATTTTCAAGAATTTTCCAATGCTACATCTTAACTTAACATTTTTATTCTT
 B.rupestris AATTTGAGAATTTTCAAGAATTTTCCAACGCTACATCTTAACTTAAATTTTTATCCTT
 B.lapidarius -----
 B.campestris AATTTGAGAATTTTCAAGA-----

B.lucorum GTATCATGTTAACGCATCGACTCTTTAAATACTACGAACGCATATATGGTACACGTGCAC
 B.terrestris GTATCAGGTTAACGCATCAACCCCTTTAAATACTACGAACGCATATATGGTACACGTGCAC
 B.t.xanthopus GTATCAGGTTAACGCATCAACCCCTTTAAATACTACGAACGCATATATGGTACACGTGCAC
 B.pascuorum GTGTCACATTAACGCATCAACCCCTTTGAATACTATGAACGCATATATGGTACACGTGCAC
 B.ruderatus GTGTCACGTTAACGCTTCAACCCCTTTAAATGCTATGAACGCATATACGGTACACGTGCAC
 B.bohemicus ATATCAGGTTAACGCTTCAACCCCTTTAAATACTATGAACGCATATATGGTACACGTGCAC
 B.maxilloso ATATCAGGTTAACGCTTCAACCCCTTTAAATACTATGAACGCATATATGGTACACGTGCAC
 B.perezi ATATCAGGTTAACGCTTCAACCCCTTTAAATACTATGAACGCATATATGGTACACGTGCAC
 B.vestalis ATATCAGGTTAACGCTTCAACCCCTTTAAATACTATGAACGCATATATGGTACACGTGCAC
 B.rupestris ATATCAGGTTAACGCTTCAACCCCTTTAAATACTATGAACGCATATATGGTACACGTGCAC
 B.lapidarius GTATCAGGTTAACGCTTCAACCCCTTTAAATACTATAAACGCATATATGGTACACGTGCAC
 B.campestris -----ACCATACCATGGA-----

| | |
|-----------------|--|
| B. lucorum | ATGGCACATTCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. terrestris | ATGGCACATTCTATTAATATCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. t. xanthopus | ATGGCACATTCTATTAATATCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. pascuorum | ATGGAACAATCTATTAATACCATTAACAAAAAGGAGGAAATAAACAGATTAACTTCG |
| B. ruderalis | ATGAAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. bohemicus | ATGGAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. maxillosus | ATGGAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. perezi | ATGGAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. vestalis | ATGGAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGATTAACTTCG |
| B. rupestris | ATGGAACAATCTATTAATACCATTAACAAAAAGGAAGAAATAAACAGGTTAACTTCG |
| B. lapidarius | -----TTAAATACCATTAACAAAAAGGCAGAAATAAACAGATTAACTTCG |
| B. campestris | ----- |
| | |
| B. lucorum | TTCAGGAAGCTTAAATAATTCCAATAGAAATCCATTTTTTAAATCGCGTCCCTATCGCAC |
| B. terrestris | TTCAGGAAGCTTAAATAATTCCAATAGAAATCCATTTTTTAAATCGCGTCCCTATCGCAC |
| B. t. xanthopus | TTCAGGAAGCTTAAATAATTCCAATAGAAATCCATTTTTTAAATCGCGTCCCTATCGCAC |
| B. pascuorum | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTCTTAAATCCCGCCCTATTGCAC |
| B. ruderalis | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTTTTAAATCCCGTCCCTATCGCAC |
| B. bohemicus | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTTTTAAATCGCGTCCCTATCGCAC |
| B. maxillosus | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTCTTAAATCGCGTCCCTATCGCAC |
| B. perezi | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTCTTAAATCGCGTCCCTATCGCAC |
| B. vestalis | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTCTTAAATCGCGTCCCTATCGCAC |
| B. rupestris | TTCAGGAACCTTAAATAATTCCAATAGAAATCCATTTCTTAAATCGCGTCCCTATCGCAC |
| B. lapidarius | TTCAGGAAGCTTAAATAATTCCAATAGAAATCCATTTTTTA----- |
| B. campestris | ----- |
| | |
| B. lucorum | AATTCGATTAATATTAGTAGTGACATT-----TTCTCCTGCATAATAACTCCAACAGC |
| B. terrestris | AATTCGATTAATATTAGTAGTGACATT-----TTCTCCTGCATAATAACTCCAACAGC |
| B. t. xanthopus | AATTCGATTAATATTAGTAGTGACATT-----TTCTCCTGCATAATAACTCCAACAGC |
| B. pascuorum | AATTCGATTAATATTAGTAGTGACATTTTTCACATTCTCTGCATAATAACACCAACAGC |
| B. ruderalis | AATTCGATTAGTA-----TTCTCCTGCATAATAAAACCAACAGC |
| B. bohemicus | AATTCATTAATATTAGTAGTAACATTTTCAAATTCCTTGAATAATAACACCAACAGC |
| B. maxillosus | AATTCATTAATATTAGTAGTAACATTTTCAAATTCCTTGAATAATAACACCAACAGC |
| B. perezi | AATTCATTAATATTAGTAGTAACATTTTCAAATTCCTTGAATAATAACACCAACAGC |
| B. vestalis | AATTCATTAATATTAGTAGTAACATTTTCAAATTCCTTGAATAATAACACCAACAGC |
| B. rupestris | AATTCATTAATATTAGTAGTGACATTTTCAAATTCCTTACATAATAACACCAACAGC |
| B. lapidarius | -----AATATTGGTAGAGACATTTTCAAATTCCTTGCATAATAACACCAACAGC |
| B. campestris | ----- |
| | |
| B. lucorum | TTACTAATAGTCCAAATAAAATTTTCATGTGAAAATT-----CTATAAATTATCAGT |
| B. terrestris | TTACTAATAGTCCAAATAAAATTTTCATGTGAAAATT-----CTATAAATTATCAGT |
| B. t. xanthopus | TTACTAATAGTCCAAATAAAATTTTCATGTGAAAATT-----CTATAAATTATCAGT |
| B. pascuorum | TTACTAATAGACCAATAAAAGTTTCACGTGAGAGTAAATATCAGAGAGTAAATTATCAAT |
| B. ruderalis | TTACTAATAGACCAATAAAATTTTCACGTGAAAATC-----CTGTAAATTATCAAT |
| B. bohemicus | TTACCAATAGACCAATAAAAGTTTCACGTGAAAATT-----CTGTAAATTATCGAT |
| B. maxillosus | TTACTAATAGACCAATAAAAGTTTCACGTGAAAATT-----CTGTAAATTATCAAT |
| B. perezi | TTACTAATAGACCAATAAAAGTTTCACGTGAAAATT-----CTGTAAATTATCAAT |
| B. vestalis | TTACTAATAGACCAATAAAAGTTTCACGTGAAAATT-----CTGTAAATTATCAAT |
| B. rupestris | TTACTAAAGGATCAAATAAACTTTTCACGTGAAAATT-----CTGTAAATTATCAAT |
| B. lapidarius | TTACTAATAGACCAATAAAAGTTTCACGTGAAAATG-----CTGTAAATTATCAAT |
| B. campestris | ----- |
| | |
| B. lucorum | CGACCAACTATCCTATTTGACATAAAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. terrestris | CGACCAACTATCCTATTTGACATAAAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. t. xanthopus | CGACCAACTATCCTATTTGACATAAAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. pascuorum | CGATCGATTACCTATTTGACATATAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. ruderalis | CGACCGATTATCCTGTTTGATATGAAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. bohemicus | CGACCGATTATCCTATTTGACATAAAATCGTA-AAATTATAAAAGTTAAACAAAAATTCT |
| B. maxillosus | CGACCGATTATCCTATTTGACATAAAATCGTA-AAATTATAAACTGCAACAAAAATTCT |
| B. perezi | CGACCGATTATCCTATTTGACATAAAATCGTA-AAATTATAAACTGCAACAAAAATTCT |
| B. vestalis | CGACCGATTATCCTATTTGACATAAAATCGTA-AAATTATAAACTGCAACAAAAATTCT |
| B. rupestris | CCACCGATTATCCTATTTGACATAAAATCGTA-AAATTATAAACTTTAAACGAAAATTCT |
| B. lapidarius | CGACCGATTATCCTATTTGACATAAAATCGTACAAGTTATAAACATTA----- |
| B. campestris | ----- |
| | |
| B. lucorum | AAAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATTGAATATTATTCACAAC |
| B. terrestris | AAAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATTGAATATTATTCACAAC |
| B. t. xanthopus | AAAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATTGAATATTATTCACAAC |
| B. pascuorum | AGAAATTGTAGCTTGCAATTTGCTGAAATTACTAGTTAATGGAATATTATTCAGAAC |
| B. ruderalis | AAAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. bohemicus | AGAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. maxillosus | AGAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. perezi | AGAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. vestalis | AGAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. rupestris | AAAAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCAGAAC |
| B. lapidarius | --AAATTGTAGCTTGCAATTTGCTGAAGTTACTAGTTAATGGAATATTATTCACATC |
| B. campestris | ----- |

| | |
|-----------------|--|
| B. lucorum | TGCAATCAAGTATCGATATTCGAAACTTCCCTTACAGCTATCATATGAATCA----- |
| B. terrestris | TGCAATCAAGTATCGATATTCGAAACTTCCCTTACAGCTATCATATGAATCA----- |
| B. t. xanthopus | TGCAATCAAGTATCGATATTCGAAACTTCCCTTACAGCTATCATATGAATCA----- |
| B. pascuorum | TGCCATCAAGTATCGATATTCGAAACTTCTCTTACAGCCACCATAACGTAAACTTTTCCC |
| B. ruderatus | TGCTATCGAGTATCGATATTCGAAACTTCCCTTACAGCCGTCATATCGTAAACTTTTCCC |
| B. bohemicus | TGCCATCAAGTATCGATATTTGAAACTTCCCTTGCAGCCGTCATATCGTAAACTTTTCCCC |
| B. maxillosus | TGCCATCAAGTATCGATATTCGAAACTTCCCTTACAGCCGTCATATCGTAAACTTTTCCCC |
| B. perezi | TGCCATCAAGTATCGATATTCGAAACTTCCCTTACAGCCGTCATATCGTAAACTTTTCCCC |
| B. vestalis | TGCCATCAAGTATCGATATTCGAAACTTCCCTTACAGCCGTCATATCGTAAACTTTTCCCC |
| B. rupestris | TGCCATCAAGTATCGATATTCGAAACTTCCCTTGCAGCCGTCGATATCGTAAACTTTTCCCC |
| B. lapidarius | TGCAATCAAGTATCGATATTTGAAACTTCCCTTACAGCCATCATATCGTAAACTTTTCCCC |
| B. campestris | -----ATTGAAACTTCCCTTACAGCCGTCATATCGTAAACTTTTCCCC |

*** ***** **.**** .*.**: .:..*

| | |
|-----------------|---|
| B. lucorum | ----- |
| B. terrestris | ----- |
| B. t. xanthopus | ----- |
| B. pascuorum | TTCTGTCGGAATCCC-TACCCACTATTTCTTTGTTTCCAATTCCTTTCTCTATTGATCCCG |
| B. ruderatus | TTC-TTCGAATCCC-TACCCCTCC-TCTTTTCGTTTCAAATTTCTTTTCTATCGATCTCG |
| B. bohemicus | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. maxillosus | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. perezi | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. vestalis | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. rupestris | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. lapidarius | TTCTGTCGGAATCCC-TACCCCTCCATCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCG |
| B. campestris | TTCTGTCGGAATCCC-TACTCTCCGTCCTCTTCGTTTCAAATTCCTTTTCTATCGATCCCA |

| | |
|-----------------|---|
| B. lucorum | ---ATATATCATGAAAAAGGTTCTGGTCCCGTTCTTCGTTATCTGTGAATCCATATTTT |
| B. terrestris | ---ATATATCATGAAAAAGGTTCTGGTCCCGTTCTTCGTTATCTGTGAATCCATATTTT |
| B. t. xanthopus | ---ATATATCATGAAAAAGGTTCTGGTCCCGTTCTTCGTTATCTGTGAATCCATATTTT |
| B. pascuorum | AATATGTTCCATGAGAAAAAGGTTCTAGTCCCATTCTTCGTTAACTGTAAATCCATATTTT |
| B. ruderatus | AATATGTTCCATGAGTAAAGGTTCTAGTCCCATTCTACGTTAACTGTGAATCCATATTTT |
| B. bohemicus | AATATGTTCCATTAGAAAAAGGTTCTATTACCGTTCTTCGTTAACTATGAATCCATATTTT |
| B. maxillosus | AATATGTTCCATTAGAAAAAGGTTCTATTCCCGTTCTTCGTTAACTATGAATCCATATTTT |
| B. perezi | AATATGTTCCATTAGAAAAAGGTTCTATTCCCGTTCTTCGTTAACTATGAATCCATATTTT |
| B. vestalis | AATATGTTCCATTAGAAAAAGGTTCTATTCCCGTTCTTCGTTAACTATGAATCCATATTTT |
| B. rupestris | AATATGTTCCATGAGAAAAAGGTTCTATTCCCGTTCTTCGTTAACTATGAATCCATATTTT |
| B. lapidarius | AATATGTTCCATGAGAAAAAGGTTCTAGTCCCATTCTTCGTTAGCTGTGAATCCATATTTT |
| B. campestris | AATATGTTCCATGAGAAAAAGGTTCTATTCCCGTTCTTCGTTAACTATGAATCCATATTTT |

.*.: *** *.:**** *.** *****:***** *.*.*****

pro peptide mature peptide

| | |
|-----------------|---|
| B. lucorum | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCTGACCCGCAAGGGTCCCTTGTTA |
| B. terrestris | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCTGACCCGCAAGGGTCCCTTGTTA |
| B. t. xanthopus | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCTGACCCGCAAGGGTCCCTTGTTA |
| B. pascuorum | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCGTCGTTA |
| B. ruderatus | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCCTCGTTA |
| B. bohemicus | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGATACGCGGACCCCAAGGGTCCGTCGTTA |
| B. maxillosus | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCGTCGTTA |
| B. perezi | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCGTCGTTA |
| B. vestalis | TTTCTTCTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCGTCGTTA |
| B. rupestris | TTTCTTTCACAGGGCTCTCCTCGTTTTTCGACGATACGCGGACCCCAAGGGTCCGTCGTTA |
| B. lapidarius | TTTCTTTTCAGGGCTCTCCTCGTTTTTCGACGACACGCGCGACCCCAAGGGTCCCTCGTTA |
| B. campestris | TTTCTTTTCAGGGCTCTCCTCGTTTTTCGACGATACGCGGACCCCAAGGGTCCATCGTTA |

***** :*****.***** ***** ***** ***** *


| | |
|-----------------|---|
| B. lucorum | TCGATGGAAAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. terrestris | TCGATGGAAAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. t. xanthopus | TCGATGGAAAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. pascuorum | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. ruderatus | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. bohemicus | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. maxillosus | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. perezi | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. vestalis | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. rupestris | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. lapidarius | TTCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |
| B. campestris | TCCAGGGACAGAAACCATTGAGCGGACCGGATCGTCGCCATCCTTGGACGTCGATTATC |

* * ***.***** ***** ***** *****

| | |
|-----------------|--|
| B. lucorum | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. terrestris | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. t. xanthopus | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. pascuorum | ATCAACGCGTCTATGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. ruderatus | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. bohemicus | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. maxillosus | ATCAACGCGTCTACGACAGAAACGGAGTGAATGCGGACGCGTACGGTGGACTGAATATTC |
| B. perezi | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. vestalis | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. rupestris | GTCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |
| B. lapidarius | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC |

| | |
|---------------|---|
| B.campestris | ATCAACGCGTCTACGACAGAAACGGAGTGAACGCGGACGCGTACGGTGGACTGAATATTC
***** |
| B.lucorum | ----- |
| B.terrestris | GCCCAGGACAGCCTGCTCAACCAC----- |
| B.t.xanthopus | GCCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATCCAGCGTGAATACAAGAATG |
| B.pascuorum | GTCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATCGGGCGTGAATACAAGAATG |
| B.ruderatus | GTCCAGGACAGCCAGCTCAACCACA----- |
| B.bohemicus | GTCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATCGGGCGTGAATACAAGAAT- |
| B.maxillosus | GTCCAGGACAGC----- |
| B.perezi | GTCCAGGACAGCC----- |
| B.vestalis | GTCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATCGGGCGTGAATACAAGAAT- |
| B.rupestris | GTCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATC----- |
| B.lapidarius | GTCCAGGACAGCCTGCTCAACCACACTTGGGTATCCAAGTCGGACGTGAATACAAGAATG |
| B.campestris | GTCCAGGACAGCCTGCTCAACCACACTTGGGTGTCCAAATCGGGCGTGAATACAAGAATG |
| B.lucorum | -- |
| B.terrestris | -- |
| B.t.xanthopus | G- |
| B.pascuorum | GC |
| B.ruderatus | -- |
| B.bohemicus | -- |
| B.maxillosus | -- |
| B.perezi | -- |
| B.vestalis | -- |
| B.rupestris | -- |
| B.lapidarius | G- |
| B.campestris | G- |

Figure S3: *Hymenoptaecin* sequence alignment using MAFFT version 6 (Katoh and Toh, 2010; Katoh et al., 2002) and annotated according to Choi et al., (2008)



Figures S4: Phylogenetic relationships of the six host and parasite bumblebee couples using Maximum Likelihood analysis for genetic information of all three AMP genes and different models for (A): all coding and non-coding regions (model: Tamura 3-Parameter including gamma distribution), (B): only coding regions (model: Kimura 2-Parameter) and (C): only non-coding regions (model: Tamura 3-Parameter), including bootstrap test of phylogeny (1000 replications). Different models were selected on the lowest BIC (Bayesian Information Criterion) score using Model selection implemented in MEGA v5.1 (Tamura et al., 2011). Note different scaling between A-B and C.

References:

- Choi YS, Choo YM, Lee KS, Yoon HJ, Kim I, Je YH, Sohn HD, Jin BR (2008) Cloning and expression profiling of four antibacterial peptide genes from the bumblebee *Bombus ignitus*. *Comp Biochem Physiol B Biochem Mol Biol*, 150: 141-46
- Katoh K, Toh H (2010) Parallelization of the MAFFT multiple sequence alignment program. *Bioinformatics*, 26: 1899-900
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res*, 30: 3059-66
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*, 28: 2731-39