

International Society of Chemical Ecology

19th Annual Meeting

University of Hamburg - Germany

August 3. - 7, 2002


Scientific Programme

and

Abstracts



Universität Hamburg



MARKING PHEROMONES OF BUMBLEBEES: COMPOSITION OF THE LABIAL GLAND SECRETION OF MALES OF *BOMBUS MAGNUS*.

Irena Valterová¹, Klára Urbanová¹, Pierre Rasmont² and Michaël Terzo^{2*}

¹Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague, Czech Republic

²University of Mons-Hainaut, Laboratory of Zoology, Avenue du Champs de Mars 6, B-7000 Mons, Belgium; *FNRS Scientific Research Worker
email: irena@uochb.cas.cz, michael.terzo@umh.ac.be

Bombus magnus is a bumblebee species belonging to the subgenus *Bombus sensu stricto* and closely related to *B. cryptarum* and *B. patagiatus*. It is abundant in Atlantic heaths from Portugal to Norway [1,2]. Characters given by Rasmont to separate *B. magnus*, *B. cryptarum*, *B. terrestris* and *B. lucorum* show a great variability in coloration and a great homogeneity of genitalia which make them difficult to identify [2]. Because of this variability Williams [3] does not recognise *B. magnus* as a good species. As application of the Paterson species recognition concept [4], the composition of the labial gland secretion, used by males for scent-marking their patrolling nuptial behaviour, seems to be the most reliable trait for recognising the validity of the bumblebees species [5].

Main component of the labial gland secretion of *B. magnus* is ethyl dodecanoate (23 %) as it is the case also for *B. patagiatus* [5] and *B. cryptarum* [6]. This is also a major or minor characteristic component in all known sexual pheromones of other species of the subgenus *Bombus* [5-7]. Unsaturated C₁₈ alcohols and ethyl esters of the corresponding acids are found in *B. magnus* and *B. lucorum* [7], but they are not reported in *B. cryptarum* [6]. Beside the main component, many other lower-abundant components are in common for *B. magnus* and *B. cryptarum*. However, their proportions are different. Another characteristic of *B. magnus* sexual pheromone is that it contains small amount (0.4 %) of geranylcitronellol and other terpene derivatives. The same is true for *B. lucorum* (0.3 % of isoprenoids). However, *B. terrestris* is the only other species of the subgenus *Bombus* which contains large amounts of isoprenoids (2,3-dihydrofarnesal 5 %, 2,3-dihydrofarnesol 18 %, 2,3-dihydrofarnesyl acetate 2 %, geranylcitronellal 4 %, geranylcitronellol 8 %, and 2,3-dihydrofarnesyl dodecanoate 6 %).

- [1] Løken, A.; Norsk Entomologisk Tidsskrift, **20**; (1973), 1-218.
- [2] Rasmont, P.; Spixiana, München, **7**; (1984), 135-160.
- [3] Williams, P.H.; Bulletin of the Natural History Museum, Entomology Series, **67**; (1998), 79-152.
- [4] Paterson, H.E.; Transvaal Museum Monograph, **4**; (1985), 21-29.
- [5] Bergström, G.; in Les colloques de l'INRA, *Les Pheromones sexuelles et les mediateurs chimiques chez les insectes: utilisation en lutte integree*, Colmar, 25-27 novembre, (1980), 35-47.
- [6] Bertsch, A.; Entomol. Gener, **22**; (1997), 129-145.
- [7] Urbanová, K., Valterová, I., Hovorka, O., Kindl, J.; Eur. J. Entomol., **98**; (2001), 111-115.